

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
15 November 2001 (15.11.2001)

PCT

(10) International Publication Number
WO 01/85912 A2

- (51) International Patent Classification⁷: C12N MCGRATH, Garth, Joseph [US/US]; Exelixis, Inc., 170 Harbor Way, P.O. Box 511, South San Francisco, CA 94083-0511 (US).
- (21) International Application Number: PCT/US01/14648
- (22) International Filing Date: 3 May 2001 (03.05.2001) (74) Agent: OSMAN, Richard, Aron; Science & Technology Law Group, 75 Denise Drive, Hillborough, CA 94010 (US).
- (25) Filing Language: English
- (26) Publication Language: English (81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.
- (30) Priority Data: 09/568,942 5 May 2000 (05.05.2000) US
- (71) Applicant (*for all designated States except US*): EX-ELIXIS, INC. [US/US]; 170 Harbor Way, P.O. Box 511, South San Francisco, CA 94083-0511 (US).
- (72) Inventors; and
- (75) Inventors/Applicants (*for US only*): CURTIS, Daniel, Tim [US/US]; Exelixis, Inc., 170 Harbor Way, P.O. Box 511, South San Francisco, CA 94083-0511 (US). FRANCIS, George, Ross [US/US]; Exelixis, Inc., 170 Harbor Way, P.O. Box 511, South San Francisco, CA 94083-0511 (US). ELLIS, Michael, Christopher [US/US]; Exelixis, Inc., 170 Harbor Way, P.O. Box 511, South San Francisco, CA 94083-0511 (US). RUDDY, David, Andrew [US/US]; Exelixis, Inc., 170 Harbor Way, P.O. Box 511, South San Francisco, CA 94083-0511 (US). NICOLL, Sharmon, Monique [US/US]; Exelixis, Inc., 170 Harbor Way, P.O. Box 511, South San Francisco, CA 94083-0511 (US).
- (84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).
- Published:
— without international search report and to be republished upon receipt of that report
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

(54) Title: PRESENILIN ENHANCERS

(57) Abstract: The invention provides methods and compositions relating to pen polypeptides having pen-specific structure and activity, related polynucleotides and modulators of pen function. The invention provides isolated pen hybridization probes and primers capable of specifically hybridizing with natural pen genes, pen-specific binding agents such as specific antibodies, and methods of making and using the subject compositions in diagnosis (e.g. genetic hybridization screens for pen transcripts), therapy (e.g. pen inhibitors to modulate APP processing) and in the biopharmaceutical industry (e.g. as immunogens, reagents for screening chemical libraries for lead pharmacological agents, etc.).

WO 01/85912 A2

Presenilin Enhancers

Inventors: Daniel Tim Curtis, George Ross Francis, Garth Joseph McGrath, Sharmon
Monique Nicoll, David Andrew Ruddy & Michael Christopher Ellis

5 Assignee: Exelixis Pharmaceuticals

INTRODUCTION

Field of the Invention

The field of this invention is proteins which modulate presenilin function.

10

Background

Alzheimer's disease is a degenerative disorder of the central nervous system which causes memory impairment and cognitive loss during mid to late life. The disease is characterized by two primary pathological features, extracellular amyloid plaques in the brain, and intra-neuronal neurofibrillary tangles. These lesions inhibit neuronal and glial cell function, and lead to synaptic loss and dementia. Both early and late onset forms of the disease have been shown to have genetic components, and four genes have been definitively associated with increased risk for AD: APP, PS1, PS2 and ApoE. These genes are functionally linked by their roles in the production, transport, and/or elimination of, amyloid- β ($A\beta$), the primary constituent of Alzheimer's amyloid plaques (reviewed in Selkoe, D. 1999, Nature 399 supp: A23).

15

20

25

30

Alzheimer's amyloid plaques are comprised largely of the 40-42 amino acid peptide $A\beta$ (Glenner, G. G., and Wong, C. W., 1984 Biochem. Biophys. Res. Commun. 122:1131). $A\beta$ is derived by proteolytic cleavage from the b-Amyloid Precursor Protein, or β APP (Kang J. et al. 1987, Nature 325:733). Three secretase activities cleave APP to generate the $A\beta$ peptide or a shorter, alternative cleavage product called p3. β -secretase generates the N-terminus of $A\beta$, while α -secretase cleaves internal to $A\beta$ sequences to generate the N-terminus of p3. γ -secretase cleaves the C-terminal β and α secretase products of APP to generate the heterogeneous C-terminal ends of $A\beta$ and p3. APP mutations found in familial Alzheimer's disease (FAD) pedigrees are clustered around the three secretase cleavage sites

(Goate, A., et al. 1991, Nature 349:704; Murrell, J., et al. 1991, Science 254: 97; Chartier-Harlin et al. 1991, Nature 353: 844; Mullan, M. et al. 1992, Nature Genet. 1: 345; Levy, E. et al., 1990, Science 248: 1124; Hendriks, L. et al. 1992, Nature Genet. 1:218) and they each increases total A β (A β 42 + A β 40) or increases the A β 42/40 ratio. Since A β 42 precipitates more readily in vitro and is the primary component of early forms of amyloid deposits called diffuse plaques, it has been postulated that increased systemic A β 42 could lead to earlier formation of plaque, and earlier onset of AD.

Family studies identified two other genes, presenilin-1 (PS1) and presenilin-2 (PS2), associated with dominantly inherited, early onset AD, (Sherrington, R. et al. 1995, Nature 375: 754; Levy-Lahad, E. et al. 1995, Science 269: 973; Rogaev., E. I. et al. 1995, Nature 376: 775). These proteins are similar to each other in sequence and encode polytopic membrane proteins with 8 transmembrane segments. Studies in FAD human cell lines, in transfected cells, and in transgenic mice have demonstrated that the PS FAD mutations cause a change in the processing pattern of APP, resulting in an increased ratio of A β 42/40 (Scheuner, D. et al. 1996, Nat. Med. 2: 864; Citron, M. et al. 1997, Nat. Med. 3:67; Borchelt, D. et al. 1996, Neuron 17: 1005; Duff, K. et al. 1996, Nature 383: 710; Tomita, T. et al. 1997, PNAS 94:2025). Studies on PS1 knockout mice demonstrated that loss of PS1 function leads to reduction in A β production due to a reduction of γ -secretase activity (De Strooper, B. et al. 1998, Nature 391: 387). Presenilin function is thus implicated in the activity of γ -secretase in two ways: missense mutations alter γ -secretase cleavage specificity, while loss of presenilin activity leads to loss of γ -secretase activity.

Inhibition of presenilin activity decreases A β production and is thus a potentially useful therapeutic approach to Alzheimer's disease. However, despite the functional link to γ -secretase activity and the generation of A β , the biochemical nature of PS activity is poorly understood. Various functions have been proposed, including action in the ER and/or Golgi complex as a chaperone for APP, Notch, and/or γ -secretase (Thinakaran, G. et al. 1998, Neurobiol. Dis. 4: 438), activity as a novel aspartyl protease, i.e. as γ -secretase itself (Wolfe, M. S. et al. 1999, Nature 398: 513), and potential roles in the response to oxidative stress and apoptosis (Wolozin, B. et al. 1996, Science 274:1710; Vito, P. et al. 1997, J. Biol. Chem 272: 28315; Guo, Q., et al. 1997, J. Neurosci. 17: 4212). The absence of a clear functional assay increases the difficulty of designing useful small molecule therapeutics targeted at presenilin.

An alternative strategy to targeting presenilin is to discover additional proteins which act together with presenilins in the pathway of γ -secretase and A β production and which might be more amenable to drug development. One useful method for the discovery of such novel targets is to perform genetic screens in model organisms such as *Drosophila* and *C. elegans* for genes that interact with presenilins.

5 Invertebrate orthologues of the PS genes have been identified by both sequence searches and genetic screens. The *C. elegans* genome contains three presenilin genes, *sel-12* (suppressor and/or enhancer of *lin-12*; Levitan, D. et al. 1995, *Nature* 377:351), *hop-1* (homolog of presenilin; Li, X. et al, 1997, *PNAS* 94:12204) and *spe-4* (spermatogenesis defective; L'Hernault et al., 1992, *J. Cell Biol.* 119:55). *sel-12*, *hop-1* and *spe-4* have 48, 35
10 and 23% sequence similarity, respectively, to PS1 and 2. *sel-12* and *hop-1* have overlapping functions in several tissues (see below), while *spe-4* appears to perform an independent function in the male germ line. Rescue experiments using transgenes have shown that human PS1 and PS2 can rescue phenotypes caused by loss of *sel-12*, demonstrating that presenilin function has been conserved from nematodes to mammals (Levitan, D. et al. 1996, *Nature*
15 377:351; Baumeister, R. et al. 1997, *Genes Function* 1: 149).

Sel-12 was identified genetically as a suppressor of an activated allele of the Notch gene *lin-12*. This discovery established a functional link between presenilin activity and activity of the Notch signaling pathway. In vivo experiments in mice (Herreman, A. et al. 1999, *PNAS* 96:11872), *Drosophila* (Struhl, G. et al. 1999, *Nature* 398: 522; Ye, Y. et al.
20 1999 *Nature* 398:525) and *C. elegans* (Li, X. et al, 1997, *PNAS* 94:12204; Westlund, B. et al. 1999, *PNAS* 96:2497) have demonstrated that the phenotype of complete loss of presenilin activity corresponds very well with the complete elimination of Notch signaling in the organism, suggesting that presenilins are absolutely required for Notch signaling activity. Notch receptors are single pass transmembrane proteins present at the cell surface that
25 mediate cell-cell signaling events critical to the differentiation of many embryonic and adult tissues in invertebrates and vertebrates. Signaling involves ligand-dependent cleavage of Notch at the inner face of the transmembrane segment, and subsequent nuclear translocation of the C-terminal domain. Analysis of Notch processing in cell culture and in vivo has further demonstrated that presenilins are required for the ligand dependent cleavage event that
30 releases the Notch intracellular domain from the transmembrane domain (Struhl, G. et al.

1999, Nature 398: 522; De Strooper, B. et al. 1999 Nature 398: 518). The parallel requirement for presenilin in both the Notch and APP cleavages suggests that the Notch signaling pathway could be a useful surrogate assay in place of Ab production in screens for presenilin pathway genes.

5 Mutations in the *C. elegans* presenilins *sel-12* and *hop-1* result in phenotypes associated with defective signaling by the *C. elegans* Notch receptors *lin-12* and *glp-1*. Loss of *hop-1* alone results in no obvious phenotypes. Loss of *sel-12* results in a strong egg-laying defective phenotype and vulval defects reminiscent of *lin-12* mutations. Loss of both *sel-12* and *hop-1* produces more severe Notch phenotypes than seen in *sel-12* alone. The specific phenotypes observed in the *sel-12; hop-1* double mutants depends on whether these worms
10 inherit maternal wild type presenilin activity. When maternally provided *sel-12+* activity is present, the double mutant displays a novel egg-laying defective phenotype and all progeny arrest during embryogenesis with *glp-1*-like developmental defects. In the absence of maternal *sel-12+* activity the double mutant exhibits a stronger phenotype of sterility with germline proliferation defects characteristic of *glp-1* mutants. Together, this set of properties
15 indicates that *sel-12* and *hop-1* are partially redundant and act coordinately to promote signaling by the two *C. elegans* Notch receptors.

The partial redundancy between *sel-12* and *hop-1* activities made it possible to look for enhancers of *sel-12* loss of function alleles that would produce a phenotype equivalent to the *sel-12;hop-1* double mutant. This enhancer screen identified two new genes which were
20 named *pen-1* and *2* (*pen* = presenilin enhancer) and which are required for presenilin function. Based on the phenotypes of the *pen* genes, we have identified a third presenilin enhancer gene, *aph-2*. The *pen-1*, *pen-2* and *aph-2* gene sequences identify orthologous genes in humans and other animals, including *pen-1B*. These genes and the processes they regulate are targets for the development of therapeutics for the treatment of Alzheimer's disease.

25 Relevant Art

Sequences related to a human *pen-1* are found, inter alia, in WO9855508, WO9855508, WO9906554 and in Unigene CGI-78 (GI#6911522 and GI#4929623)

Sequences related to a human *pen-2* are found, inter alia, in AD000671 (genomic) and
30 GI#3601371 (cDNA).

Sequences related to a human Aph-2 are found inter alia, in WO 9845435, WO 9845436, WO 9300353 and (KIAA0253, DNA GI1665772, protein GI 1665773).

Numerous ESTs were found in public databases containing pieces of the natural human pen-1B sequence disclosed herein, including ns43g08.s1 (GI# 2874520, not annotated) and ESTs of Unigen contig Hs.42954 (53% similar to pen-1 (CGI-78)), including:

5 AI538204 (IMAGE:2189986); AA808355 (IMAGE:1334417); N21153 (IMAGE:264868); AI204164 (IMAGE:1734840); AI001990 (IMAGE:1619191); AA578718 (IMAGE:953241); AA887975 (IMAGE:1160119); AI004282 (IMAGE:1626004); AI188040 (IMAGE:1738954); AI192033 (IMAGE:1738659); AI005113 (IMAGE:1626277); AW118908 (IMAGE:2605631); AI760754 (IMAGE:2398349); AA805770 (IMAGE:1186430); AA805757 (IMAGE:1186406); AW182071 (IMAGE:2662428); AA805773 (IMAGE:1186436); AI301191 (IMAGE:1897253); AA976455 (IMAGE:1589895); and N31710 (IMAGE:271292).

SUMMARY OF THE INVENTION

15 The invention provides methods, compositions and systems relating to presenilin enhancer proteins (pens), including methods for modulating (e.g. enhancing or inhibiting) and detecting presenilin-pen interactions. In a particular embodiment, the method provides for specifically detecting a stress that alters a functional interaction of a presenilin enhancer (pen) with upstream or downstream Notch or APP processing by: (i) introducing a predetermined

20 stress into a system which provides a functional interaction of a pen with Notch or APP processing, whereby the system provides a stress-biased interaction of the pen with Notch or APP processing, wherein the absence of the stress, the system provides unbiased interaction of the pen with Notch or APP processing; and (ii) detecting the stress-biased interaction of the pen with Notch or APP processing, wherein a difference between the stress-biased and

25 unbiased interactions indicates that the stress alters the interaction of the pen with Notch or APP processing.

The system may be a viable cell expressing the pen wherein the pen expression is determined to be non-natural or pathogenic, or an in vitro, cell-free mixture comprising a determined amount of the pen. A wide variety of embodiments are encompassed; for

30 example, wherein the system is the viable cell, in situ or in vitro, and the stress is a

pharmacologically active agent or a deficiency in functional expression of the pen, such as by virtue of genomic disruption of otherwise endogenous alleles encoding the pen or coexpression of a polynucleotide comprising a sequence antisense of an endogenous allele encoding the pen. Alternatively, the system may be the in vitro, cell-free mixture and the stress is a pharmacologically active agent. The stress-biased interaction of the pen with Notch or APP processing may be detected by any convenient means or marker, such as detecting an indication of Alzheimer's disease, a transcriptional reporter of notch, generation of a downstream product such as A β or a structural alteration in the pen, such as with a specific antibody.

The invention provides a variety of other methods and compositions relating to pen polypeptides having pen-specific structure and activity, related polynucleotides and modulators of pen function. The pen polypeptides may be recombinantly produced from transformed host cells from the subject pen polypeptide encoding nucleic acids or purified from natural sources such as mammalian cells. The invention provides isolated pen hybridization probes and primers capable of specifically hybridizing with natural pen genes, pen-specific binding agents such as specific antibodies, agonists and antagonists, and methods of making and using the subject compositions in diagnosis (e.g. genetic hybridization screens for pen transcripts), therapy (e.g. pen inhibitors to modulate A β production) and in the biopharmaceutical industry (e.g. as immunogens, reagents for isolating natural pen genes and transcripts, reagents for screening chemical libraries for lead pharmacological agents, etc.). In a particular aspect, the pen methods and compositions relate to pen-1B polypeptides.

DETAILED DESCRIPTION OF THE INVENTION

The invention provides methods, compositions and systems relating to presenilin enhancer proteins (pens), including methods for modulating (e.g. enhancing or inhibiting) and/or detecting an interaction between a pen and Notch or APP processing. In a particular embodiment, the method provides for specifically detecting a stress that alters a functional interaction of a presenilin enhancer (pen) with Notch or APP processing.

The pen is independently selected from a pen-1, pen-1B, pen-2 and Aph-2 polypeptide. These names are used generically to refer to polypeptides which comprise a

disclosed parental sequence, comprise specified fragments thereof, or have sequence similarity to a disclosed parental sequence, wherein the sequence similarity is at least 40%, preferably at least 60%, more preferably at least 80%, more preferably at least 90%, more preferably at least 95%, most preferably 100%, and specifically bind a specifically disclosed presenilin or corresponding parental sequence pen-specific antibody, as measured in one or more of the disclosed interaction assays. The polypeptides comprise, and the similarity or identity extends over at least 10, preferably at least 15, more preferably at least 25, more preferably at least 35, more preferably at least 50 contiguous residues and most preferably over the entire polypeptide and/or parental pen sequence.

10 Table 1. Parental pen Polypeptides

Parental pen	Natural Source	SEQ ID NO	% identity to human parental pen by BLAST
pen-1	C. elegans	(SEQ ID NO:1)	28.7
	D. melanogaster	(SEQ ID NO:2)	45.4
	H. Virescens	(SEQ ID NO:3)	50
	mouse	(SEQ ID NO:4)	92.8
	human	(SEQ ID NO:5)	100
pen-1B	human	(SEQ ID NO:6)	51(identity to human parental pen-1)
pen-2	C. elegans	(SEQ ID NO:7)	42.6
	D. melanogaster	(SEQ ID NO:8)	60.4
	rat	(SEQ ID NO:9)	96
	mouse	(SEQ ID NO:10)	96
	cow	(SEQ ID NO:11)	95
	human	(SEQ ID NO:12)	100
Aph-2	C. elegans	(SEQ ID NO:13)	18.9
	D. melanogaster	(SEQ ID NO:14)	29.9
	human	(SEQ ID NO:15)	100

For disclosed polymeric genes, "percent (%) sequence identity over a specified window size W" with respect to parental sequences is defined as the percentage of residues in any window of W residues in the candidate sequence that are identical with the residues in the parent sequence, after aligning the sequences and introducing gaps, if necessary, to achieve the maximum percent sequence identity. The % identity values are generated by

5 WU-BLAST-2 .0 a19 obtained from Altschul et al., J. Mol. Biol., 215: 403-410(1990); <http://blast.wustl.edu/blast/README.html>. WU-BLAST-2.0a19 which uses several search parameters, all of which are set to the default values. The HSP S and HSP S2 parameters are dynamic values and are established by the program itself depending upon the composition of the particular sequence and composition of the particular database against which the sequence

10 of interest is being searched; however, the values may be adjusted to increase sensitivity. Hence, a % sequence identity value is determined by the number of matching identical residues divided by the window size W for which the percent identity is reported. Exemplary species are readily generated by mutating the corresponding parental sequences and confirming presenilin or antibody binding. For example, pen-1B polypeptides defined by

15 SEQ ID NOS:16-25 exemplify an active (demonstrating presenilin binding) 90% genus around parental sequence SEQ ID NO:6. In particular embodiments, the pen is a natural pen, such as human, mouse, *D. melanogaster*, *H. virescens* or *C. elegans* pen-1; human, rat, mouse, cow, *D. melanogaster* or *C. elegans* pen-2; human pen-1B, and human, *D. melanogaster* or *C. elegans* Aph-2. In a particular aspect, the pen is a naturally-occurring pen

20 identifiable in a sel-12Δ (Δ means deletion allele) homozygous *C. elegans* genetic mutation enhancer screen.

The interaction between the pen and Notch or APP processing may be detected in any convenient manner that specifically assays the pen influence on the processing pathway. The assay may be constructed to monitor a downstream perturbation in product generation (e.g.

25 Aβ or Notch intracellular domain production), an intermediate pathway step (a number of intermediate Notch and APP processing pathway steps and intermediate component interactions are well documented in the art), or initiating pen - presenilin or pen - γ-secretase binding.

A wide variety of systems may be used in the methods. Detailed below are animal

30 systems stressed with mutant pen genes to provide sensitized Notch and/or APP processing pathways, which systems are used to characterize additional interacting proteins. In particular embodiments, the system comprises a cell or animal expressing both the pen and a binding target such as a presenilin or γ-secretase, an in vitro, cell-free mixture comprising a determined amount of the pen and a binding target; applications of such cells and mixtures

35 include two-hybrid, biochemical pull-down, immunoprecipitation, fluorescent polarization and solid phase binding assays. In accordance with the diversity of applicable systems, a

wide variety of stresses may be assayed or evaluated, including chemical agents, such as candidate drugs, toxins, contaminants, etc.; radiation such as ultraviolet rays and x-rays; infection such as viral or bacterial infection including cellular transformation; genetic mutations, etc.

5 The particular method used to detect the interaction of the pen polypeptide and the presenilin will depend on the nature of the assay, so long as the interaction is specifically detected. For example, as detailed below, modulation of pen mutant specific phenotypes provide readouts for genetic interaction assays. For in vitro assays, depending on if and how the pen polypeptide and/or target are labeled, the interaction readout may be measured by changes in fluorescence, optical density, gel shifts, radiation, etc. In a particular embodiment, 10 the system provides a downstream APP processing readout.

In a particular embodiment, the methods involve specifically detecting a stress that alters a physical interaction of a subject pen polypeptide with APP and/or Notch processing. In one aspect, this embodiment comprises the steps of (a) introducing a predetermined stress into a system which provides a physical interaction of a pen with a binding target, whereby 15 the system provides a stress-biased interaction of the pen and the target, wherein the absence of the stress, the system provides an unbiased interaction of the pen polypeptide and the target; and (b) detecting the stress-biased interaction of the pen polypeptide and the target, wherein a difference between the stress-biased and unbiased interactions indicates that the stress alters the interaction of the pen polypeptide and the target, wherein preferred targets include γ -secretases, presenilins, notch and/or APP substrates, and/or combinations and 20 complexes thereof.

In the latter embodiment, the presenilin is selected from a presenilin-1 (PS-1) and presenilin-2 (PS-2). These names are used generically to refer to polypeptides which comprise a disclosed parental sequence, comprises specified fragments thereof, or have 25 sequence similarity to the disclosed parental presenilin sequences, wherein the sequence similarity is at least 50%, more preferably at least 70%, more preferably at least 80%, more preferably at least 90%, more preferably at least 95% and most preferably 100%, wherein the presenilin is sufficient to provide a presenilin-specific, detectable functional interaction comparable to that provided by the corresponding parental sequence presenilin, as measured 30 in one or more of the disclosed genetic or biochemical interaction assays. The presenilins comprise, and the similarity or identity extends over at least 10, preferably at least 15, more preferably at least 25, more preferably at least 35, more preferably at least 50 contiguous residues and most preferably over the entire presenilin or parental sequence. The parental presenilin is selected from a natural sequence presenilin 1 (such as human, mouse, chicken

and xenopus sequences) and presenilin 2 (such as human, mouse and xenopus sequences), which are known in the art and accessible from public genetic depositories such as Genbank.

The compositions of the invention, useful in the subject methods, include the subject pen polypeptides and mixtures comprising predetermined amounts of a disclosed pen and presenilin polypeptides, particularly wherein one, preferably both of these components are isolated and mixtures consisting essentially of both components, i.e. wherein other components of the mixture (except for an assayed stress) do not significantly influence the interaction of these two components. Other aspects of the invention include nucleic acids encoding the disclosed pen polypeptides, antibodies which specifically bind them, and methods of use.

Subject polypeptides consisting of the disclosed parental sequences or fragments thereof are isolated, i.e. encompass pen polypeptides covalently joined to a non-natural or heterologous component, such as a non-natural amino acid or amino acid sequence or a natural amino acid or sequence other than that which the polypeptide is joined to in a natural protein, are preferably in solution, and preferably constitute at least about 0.5%, and more preferably at least about 5% by weight of the total polypeptide in a given sample and pure polypeptides constitute at least about 90%, and preferably at least about 99% by weight of the total polypeptide in a given sample, as are preferred subject polypeptides comprising other than parental sequence. The polypeptides may be covalently or noncovalently part of a larger complex, such as larger polypeptides and/or various conjugates, etc. The polypeptides may be synthesized, produced by recombinant technology, or purified from mammalian, preferably human cells. A wide variety of molecular and biochemical methods are available for biochemical synthesis, molecular expression and purification of the subject compositions, see e.g. Molecular Cloning, A Laboratory Manual (Sambrook, *et al.* Cold Spring Harbor Laboratory), Current Protocols in Molecular Biology (Eds. Ausubel, *et al.*, Greene Publ. Assoc., Wiley-Interscience, NY) or that are otherwise known in the art.

The pen encompassing fragments comprise at least 10, preferably at least 15, more preferably at least 25, more preferably at least 35, most preferably at least 50 consecutive residues of a corresponding disclosed parental pen sequence. Pen polypeptides provide corresponding pen specific function, such as interacting with a component of a natural notch or APP processing pathway, especially presenilin binding or binding inhibitory activity as shown in one or more binding assays as described herein, and/or pen specific antibody binding or binding inhibitory activity, particularly as measured in a disclosed binding assay.

Pen-specific function may be determined by convenient *in vitro*, cell-based, or *in vivo* assays, e.g. binding assays. The term binding assay is used generically to encompass any assay, including *in vitro*, cell-culture or animal-based assays (e.g. using gene therapy

techniques or with transgenics), etc. where the molecular interaction of a pen polypeptide with a specific binding target is evaluated. The binding target may be a natural intracellular binding target such as a presenilin, a pen regulating protein or other regulator that directly modulates pen activity or its localization; or non-natural binding target such as a specific immune protein such as an antibody, or a pen specific agent such as those identified in screening assays such as described below. Pen-binding specificity may be assayed by APP processing (e.g. ability of the subject polypeptides to function as negative effectors in pen-expressing cells), by binding equilibrium constants (usually at least about 10^7 M^{-1} , preferably at least about 10^8 M^{-1} , more preferably at least about 10^9 M^{-1}), by immunogenicity (e.g. ability to elicit pen specific antibody in a heterologous host such as a mouse, rat, goat or rabbit), etc.

In a particular embodiment, the subject polypeptides provide pen-specific antigens and/or immunogens, especially when coupled to carrier proteins. For example, the subject polypeptides are covalently coupled to keyhole limpet antigen (KLH) and the conjugate is emulsified in Freund's complete adjuvant. Laboratory rabbits are immunized according to conventional protocol and bled. The presence of pen-specific antibodies is assayed by solid phase immunosorbant assays using immobilized corresponding pen polypeptides, see, e.g. Table 2.

Table 2. Immunogenic pen-1B polypeptides eliciting pen-1B-specific rabbit polyclonal antibody: pen-1B polypeptide-KLH conjugates immunized per protocol described above.

<u>pen-1B</u>	<u>Immuno-</u>	<u>pen-1B</u>	<u>Immuno-</u>
<u>Polypeptide Sequence</u>	<u>genicity</u>	<u>Polypeptide Sequence</u>	<u>genicity</u>
SEQ ID NO:6, res 1-14	+++	SEQ ID NO:6, res 115-126	+++
SEQ ID NO:6, res 6-15	+++	SEQ ID NO:6, res 130-140	+++
SEQ ID NO:6, res 10-20	+++	SEQ ID NO:6, res 139-151	+++
SEQ ID NO:6, res 25-46	+++	SEQ ID NO:6, res 166-182	+++
SEQ ID NO:6, res 62-71	+++	SEQ ID NO:6, res 184-198	+++
SEQ ID NO:6, res 67-76	+++	SEQ ID NO:6, res 214-232	+++
SEQ ID NO:6, res 72-95	+++	SEQ ID NO:6, res 246-257	+++

The subject pen polypeptides also encompass minor deletion mutants, including N-, and/or C-terminal truncations, of the parental pen polypeptides. Such deletion mutants are readily screened for pen competitive or dominant negative activity. Exemplary active

deletion mutants for pen-1B include polypeptides comprising an amino acid sequence selected from the group consisting of: SEQ ID NO:6, residues 1-254; SEQ ID NO:6, residues 4-255; SEQ ID NO:6, residues 9-257; and SEQ ID NO:6, residues 2-255.

The invention provides binding agents specific to the claimed pen-1B polypeptides, including natural intracellular binding targets, etc., methods of identifying and making such agents, and their use in diagnosis, therapy and pharmaceutical development. For example, specific binding agents are useful in a variety of diagnostic and therapeutic applications, especially where disease or disease prognosis is associated with unoptimized utilization of a pathway involving pen, e.g. APP processing. Novel pen-specific binding agents include pen-specific receptors, such as somatically recombined polypeptide receptors like specific antibodies or T-cell antigen receptors (see, e.g. Harlow and Lane (1988) *Antibodies, A Laboratory Manual*, Cold Spring Harbor Laboratory) and other natural intracellular binding agents identified with assays such as one-, two- and three-hybrid screens, non-natural intracellular binding agents identified in screens of chemical libraries such as in vitro, cell-based and animal-based binding assays described herein, or otherwise known to those of skill in the art, etc. Agents of particular interest modulate pen function, e.g. pen-dependent Notch or APP processing, and include dominant negative deletion mutants, etc. Accordingly, the invention also provides methods for modulating APP processing in a cell comprising the step of modulating pen activity, e.g. by contacting the cell with a modulator of a resident pen, a dominant negative pen deletion mutant, or pen polynucleotide (below).

In addition to direct synthesis, the subject polypeptides can also be expressed in cell and cell-free systems (e.g. Jermutus L, et al., *Curr Opin Biotechnol.* 1998 Oct;9(5):534-48) from encoding polynucleotides, such as the corresponding parent polynucleotides or naturally-encoding polynucleotides isolated with degenerate oligonucleotide primers and probes generated from the subject polypeptide sequences ("GCG" software, Genetics Computer Group, Inc, Madison WI) or polynucleotides optimized for selected expression systems made by back-translating the subject polypeptides according to computer algorithms (e.g. Holler et al. (1993) *Gene* 136, 323-328; Martin et al. (1995) *Gene* 154, 150-166). Hence, the polypeptides may be synthesized, produced by recombinant technology, or purified from cells. A wide variety of molecular and biochemical methods are available for biochemical synthesis, molecular expression and purification of the subject compositions, see e.g. *Molecular Cloning, A Laboratory Manual* (Sambrook, et al. Cold Spring Harbor Laboratory), *Current Protocols in Molecular Biology* (Eds. Ausubel, et al., Greene Publ. Assoc., Wiley-Interscience, NY) or that are otherwise known in the art.

The invention provides polynucleotides encoding the disclosed polypeptides, and pen-gene specific polynucleotides, which polynucleotides may be joined to other components

such as labels or other polynucleotide sequences (i.e. they may be part of larger sequences) and are of synthetic/non-natural sequences and/or are isolated, i.e. unaccompanied by at least some of the material with which it is associated in its natural state, preferably constituting at least about 0.5%, preferably at least about 5% by weight of total nucleic acid present in a given fraction, and usually recombinant, meaning they comprise a non-natural sequence or a natural sequence joined to nucleotide(s) other than that which it is joined to on a natural chromosome. Recombinant polynucleotides comprising natural sequence contain such sequence at a terminus, immediately flanked by (i.e. contiguous with) a sequence other than that which it is joined to on a natural chromosome, or flanked by a native flanking region fewer than 10 kb, preferably fewer than 2 kb, more preferably fewer than 500 bases, most preferably fewer than 100 bases, which is at a terminus or is immediately flanked by a sequence other than that which it is joined to on a natural chromosome. While the polynucleotides are usually RNA or DNA, it is often advantageous to use polynucleotides comprising other bases or nucleotide analogs to provide modified stability, etc. Furthermore, the terms polynucleotide and nucleic acid are used interchangeably to refer to any polymer of nucleotides, without restriction by length.

The invention also encompasses pen, particularly pen-1B gene specific polynucleotides. For example, the nucleotide sequence of a natural human transcript encoding a natural human pen-1B polypeptide is shown as SEQ ID NO:26. The term pen-1B gene specific polynucleotides is used generically to refer to polynucleotides comprising SEQ ID NO:26, comprising specified fragments of SEQ ID NO:26, or having sequence similarity to SEQ ID NO:26. Subject fragments of SEQ ID NO:26, which are useful, e.g. as hybridization probes and replication / amplification primers, comprise at least 12, preferably at least 24, more preferably at least 48, more preferably at least 96 and most preferably at least 182 contiguous nucleotides of SEQ ID NO:26.

Pen gene specific polynucleotides effect specific hybridization to the corresponding parental sequence or complement thereof; for example, all pen-1B gene specific polynucleotides effect specific hybridization to SEQ ID NO:26 or its complement. Demonstrating specific hybridization generally requires stringent conditions, for example, hybridizing in a buffer comprising 30% formamide in 5 x SSPE (0.18 M NaCl, 0.01 M NaPO₄, pH 7.7, 0.001 M EDTA) buffer at a temperature of 42°C and remaining bound when subject to washing at 42°C with 0.2 x SSPE; preferably hybridizing in a buffer comprising 50% formamide in 5 x SSPE buffer at a temperature of 42°C and remaining bound when subject to washing at 42°C with 0.2 x SSPE buffer at 42°C. Specifically hybridizing polynucleotides are readily identified in convenient gel-based assays; for example, polynucleotides comprising SEQ ID NOS:27-38 are shown to specifically hybridize with SEQ ID NO:26 under the foregoing preferred hybridization conditions.

Table 3. Exemplary pen-1B gene specific polynucleotides which hybridize with a strand of SEQ ID NO:26 under Conditions I and II.

<u>pen-1B gene specific polynucleotides</u>	<u>Specific Hybrids</u>	<u>pen-1B gene specific polynucleotides</u>	<u>Specific Hybrids</u>
SEQ ID NO:26, nucl 1-36	+	SEQ ID NO:27	+
SEQ ID NO:26, nucl 32-68	+	SEQ ID NO:28	+
SEQ ID NO:26, nucl 65-97	+	SEQ ID NO:28	+
SEQ ID NO:26, nucl 103-140	+	SEQ ID NO:30	+
SEQ ID NO:26, nucl 131-154	+	SEQ ID NO:31	+
SEQ ID NO:26, nucl 148-182	+	SEQ ID NO:32	+
SEQ ID NO:26, nucl 222-256	+	SEQ ID NO:33	+
SEQ ID NO:26, nucl 258-286	+	SEQ ID NO:34	+
SEQ ID NO:26, nucl 273-305	+	SEQ ID NO:35	+
SEQ ID NO:26, nucl 318-352	+	SEQ ID NO:36	+
SEQ ID NO:26, nucl 344-376	+	SEQ ID NO:37	+
SEQ ID NO:26, nucl 352-386	+	SEQ ID NO:38	+
SEQ ID NO:26, nucl 388-424	+		
SEQ ID NO:26, nucl 406-431	+		
SEQ ID NO:26, nucl 420-446	+		

20 The subject nucleic acids find a wide variety of applications including use as translatable transcripts, hybridization probes, PCR primers, diagnostic nucleic acids, etc.; use in detecting the presence of other pen gene specific polynucleotides and gene transcripts and in detecting or amplifying nucleic acids encoding additional pen homologs and structural analogs. For example, pen-encoding polynucleotides may be used in pen-expression vectors, generally operably linked to a heterologous promoter, and/or incorporated into recombinant host cells, e.g. for expression and screening, transgenic animals, e.g. for functional studies such as the efficacy of candidate drugs for disease associated with pen-modulated cell function, etc. In diagnosis, pen hybridization probes find use in identifying wild-type and mutant pen alleles in clinical and laboratory samples. Mutant alleles are used to generate allele-specific probes for high-throughput clinical diagnoses, e.g. for pen mutations associated

with Alzheimer's disease. In therapy, therapeutic pen polynucleotides are used to modulate cellular expression or intracellular concentration or availability of active pen.

For example, pen polynucleotides are used to modulate cellular expression or intracellular concentration or availability of active pen protein. Pen inhibitory nucleic acids are typically antisense: single-stranded sequences comprising complements of the disclosed natural pen transcript sequence. Antisense modulation of the expression of a given pen polypeptide may employ antisense nucleic acids operably linked to gene regulatory sequences. Cells are transfected with a vector comprising a pen gene specific polynucleotide sequence with a promoter sequence oriented such that transcription of the gene yields an antisense transcript capable of binding to endogenous pen encoding mRNA. Alternatively, single-stranded antisense polynucleotides that bind to genomic DNA or mRNA encoding pen polypeptide may be administered to the target cell, in or temporarily isolated from a host, at a concentration that results in a substantial reduction in expression of the targeted protein. An enhancement in pen expression is effected by introducing into the targeted cell type pen polynucleotides that increase the functional expression of the corresponding gene products. Such polynucleotides may be pen expression vectors, vectors that upregulate the functional expression of an endogenous allele, or replacement vectors for targeted modification of endogenous mutant or wild type alleles. Techniques for introducing the nucleic acids into viable cells are known in the art and include retroviral-based transfection, viral coat protein-liposome mediated transfection, etc.

The invention provides efficient methods of identifying agents, compounds or lead compounds for agents active at the level of a pen modulatable cellular function and/or pen gene expression, including transcription. A wide variety of assays for transcriptional modulators or binding agents is provided including labeled *in vitro* ligand binding assays, immunoassays, etc. The methods are amenable to automated, cost-effective high throughput screening of chemical libraries for lead compounds. Identified reagents find use in the pharmaceutical industries for animal and human trials; for example, the reagents may be derivatized and rescreened in *in vitro* and *in vivo* assays to optimize activity and minimize toxicity for pharmaceutical development.

A wide variety of assays for binding agents, i.e. screens for compounds that modulate pen interaction with a natural pen binding target are also provided. These assays employ a mixture of components including a pen polypeptide, which may be part of a fusion product with another polypeptide, e.g. a peptide tag for detection or anchoring, etc. The assay mixtures comprise a natural intracellular pen binding target. In a particular embodiment, the binding target is presenilin, or portion thereof which provides binding affinity and avidity to the subject pen polypeptide conveniently measurable in the assay and preferably comparable

to the intact presenilin. The assay mixture also comprises a candidate pharmacological agent. Candidate agents encompass numerous chemical classes, though typically they are organic compounds; preferably small organic compounds and are obtained from a wide variety of sources including libraries of synthetic or natural compounds. A variety of other reagents may also be included in the mixture. These include reagents like salts, buffers, neutral proteins, e.g. albumin, detergents, protease inhibitors, nuclease inhibitors, antimicrobial agents, etc.

The resultant mixture is incubated under conditions whereby, but for the presence of the candidate pharmacological agent, the pen polypeptide specifically binds the cellular binding target, portion or analog with a reference binding affinity. The mixture components can be added in any order that provides for the requisite bindings, and incubations may be performed at any temperature which facilitates optimal binding. Incubation periods are likewise selected for optimal binding but also minimized to facilitate rapid, high-throughput screening.

After incubation, the agent-biased binding between the pen polypeptide and one or more binding targets is detected by any convenient way. A variety of methods may be used to detect the change depending on the nature of the product and other assay components, e.g. through optical or electron density, radiative emissions, nonradiative energy transfers, etc. or indirect detection with antibody conjugates, etc. A difference in the binding affinity of the pen-1B to the target in the absence of the agent as compared with the binding affinity in the presence of the agent indicates that the agent modulates the binding of the pen to the pen binding target. A difference, as used herein, is statistically significant and preferably represents at least a 50%, more preferably at least a 90% difference.

The following experimental section and examples are offered by way of illustration and not by way of limitation.

EXAMPLES, PROTOCOLS AND EXPERIMENTAL PROCEDURES

I. High-Throughput In Vitro Fluorescence Polarization Assay

Reagents:

pen peptide (size minimized, rhodamine-labeled; final conc. = 1 - 5 nM)

PS polypeptide (final conc. = 100 - 200 nM)

Buffer: 10 mM HEPES, 10 mM NaCl, 6 mM magnesium chloride, pH 7.6

Protocol:

1. Add 90 microliters of pen peptide/PS polypeptide mixture to each well of a 96-well microtiter plate.
2. Add 10 microliters of test compound per well.
3. Shake 5 min and within 5 minutes determine amount of fluorescence polarization by using a Fluorolite FPM-2 Fluorescence Polarization Microtiter System (Dynatech Laboratories, Inc).

II. Conformational Sensor - ELISA Format Assay

Buffer and Solution Preparation:

1. 10X Assay Buffer:

100mL of 1M Hepes

300mL of 5M NaCl

20mL of 1M MgCl

Add MQ H₂O to 1L

2. Master Mix of peptide / protein

Protein: Glutathione-S-transferase/ γ -secretase polypeptide fusion protein:

final conc = 100 nM

pen peptide (size minimized, biotinylated; final conc. = 1 uM)

Add Assay Buffer and H₂O to bring to final volume: final buffer conc = 1X

3. Antibody Mix:

anti-GST, rabbit (final conc. = 1:10,000)

anti-rabbit-HRP (final conc. = 1:10,000)

Add T-TBS to bring to final volume: final buffer conc = 1X

Procedure:

1. Make 50 mL of Master Mix (see 2 above) of appropriate peptide / protein combinations (use 50 mL polypropylene tubes). Incubate for 1 hr at RT

2. Add 95 uL of Master Mix to each well of a 96-well plate**

** Reacti-Bind Streptavidin-Coated, White Polystyrene Plates (#15118B), which have been blocked by Super-Blocking Reagent from Pierce.

3. Transfer 5 uL of each test compound (stock = 60 uM) to each well of the plate

4. Incubate plate for 1hr at RT

5. While incubating, make rabbit anti-GST antibody and anti-rabbit-HRP Antibody Mix (see 3 above). Incubate on ice for 1 hr.
6. Wash plates 3X with H₂O thoroughly
7. Add 100 uL of Antibody Mix into each well of the plate
8. Incubate for 1 hr at RT
- 5 9. Wash 3X with H₂O
10. Dilute Supersignal substrate (mixed Luminol and peroxide) in 1:2 H₂O and then add 100 uL into each well
11. Shake 3-5 min. Read chemiluminescence.

10 III. High-Throughput In Vitro Binding Assay.

A. Reagents:

- Neutralite Avidin: 20 µg/ml in PBS.
- Blocking buffer: 5% BSA, 0.5% Tween 20 in PBS; 1 hour at room temperature.
- Assay Buffer: 100 mM KCl, 20 mM HEPES pH 7.6, 1 mM MgCl₂, 1% glycerol,
15 0.5% NP-40, 50 mM b-mercaptoethanol, 1 mg/ml BSA, cocktail of protease inhibitors.
- ³²P pen peptide 10x stock: 10⁻⁸ - 10⁻⁶ M "cold" pen peptide supplemented with 200,000-250,000 cpm of labeled pen peptide (Beckman counter). Place in the 4°C microfridge during screening.
- Protease inhibitor cocktail (1000X): 10 mg Trypsin Inhibitor (BMB # 109894), 10
20 mg Aprotinin (BMB # 236624), 25 mg Benzamidin (Sigma # B-6506), 25 mg Leupeptin (BMB # 1017128), 10 mg APMSF (BMB # 917575), and 2mM NaVO₃ (Sigma # S-6508) in 10 ml of PBS.

- Binding Polypeptide: 10⁻⁷ - 10⁻⁵ M biotinylated PS polypeptide in PBS.

B. Preparation of assay plates:

- 25 - Coat with 120 µl of stock N-Avidin per well overnight at 4°C.
- Wash 2 times with 200 µl PBS.
- Block with 150 µl of blocking buffer.
- Wash 2 times with 200 µl PBS.

C. Assay:

- 30 - Add 40 µl assay buffer/well.

- Add 10 μ l compound or extract.
- Add 10 μ l 33 P-pen peptide (20-25,000 cpm/0.1-10 pmoles/well = 10^{-9} - 10^{-7} M final conc).
- Shake at 25°C for 15 minutes.
- Incubate additional 45 minutes at 25°C.
- 5 - Add 40 μ M biotinylated PS polypeptide (0.1-10 pmoles/40 μ l in assay buffer)
- Incubate 1 hour at room temperature.
- Stop the reaction by washing 4 times with 200 μ M PBS.
- Add 150 μ M scintillation cocktail.
- Count in Topcount.
- 10 D. Controls for all assays (located on each plate):
 - a. Non-specific binding
 - b. Soluble (non-biotinylated PS polypeptide) at 80% inhibition.

IV. Identification of presenilin enhancer genes: natural pen-1 and pen-2.

- 15 The partial redundancy of sel-12 and hop-1 means that, for most tissues, a deletion of one or the other gene will result in only a partial loss of presenilin function. Hence, a knock-out mutation in one or the other gene provides a sensitized background for genetic screens designed to identify presenilin interacting genes. Using this reasoning, we designed several variations of genetic screens aimed at identifying genes that act in concert with
- 20 presenilins. One variation (Screen A) is to mutagenize worms homozygous for a sel-12 deletion mutation (hereinafter referred to as sel-12 Δ) and screen for enhancer mutations that, in combination with sel-12 Δ , produce phenotypes equivalent to those of the sel-12 Δ ; hop-1 Δ double mutant. Such enhancer mutations identify both 1) components that interact uniquely with hop-1 presenilin and 2) components that interact with both hop-1 and sel-12 presenilins.
- 25 As an internal control, Screen A is expected to yield loss-of-function hop-1 alleles since the screen targets the phenotypes seen in the sel-12 Δ ; hop-1 Δ double mutant. Another variation is to mutagenize a hop-1 single mutant and again screen for enhancement to the phenotypes associated with a complete presenilin.

- 30 In addition to the desired mutations that enhance presenilin defects, these screens identify mutations in known components of the glp-1 signaling pathway (e.g., glp-1/Notch receptor, lag-2/DSL ligand, lag-2/Su(H) family effector) since loss of these gene products results in glp-1 like sterility. An important distinction between presenilin enhancers and

mutations in known glp-1 pathway genes is that former result in glp-1-like sterility only in a sel-12 Δ background whereas the latter result in glp-1 sterility in both a wild-type genetic background (Austin, J. and Kimble, J., Cell (1987) 51:589-599; Lambie, E. and Kimble, J., Development 1991) 112:231-240) and a sel-12 Δ background.

We performed Screen A on a large scale, screening approximately 128,000 haploid
5 genomes after mutagenesis of a sel-12 Δ homozygous strain with ethyl methane sulfonate. The screen resulted in the isolation of the expected types of mutants, including 27 putative glp-1 alleles, 3 mutations identified as likely lag-1 or lag-2 alleles based on map position, and 8 hop-1 mutations. As expected, the putative glp-1, lag-1, and lag-2 mutations result in glp-1-like sterility in both a wild type and a sel-12 Δ genetic background; these mutations
10 therefore cause sterility independently of the presence or absence sel-12+ function. By contrast, the 8 hop-1 mutations result in a penetrant glp-1-like sterile phenotype in the absence, but not the presence, of sel-12+ activity.

In addition to the preceding, we isolated 7 mutants that, based on mapping and
complementation tests, identify two new presenilin-interacting genes. Four of these mutants
15 identify the gene pen-1 located on chromosome I and other three identify the gene pen-2 located on chromosome III. Our subsequent work with these genes indicated: 1) that the pen-1 and pen-2 enhancers alleles are loss-of-function mutations; 2) that loss of pen-1+ or pen-2+ function, in combination with a loss of sel-12+ function, has the same phenotypic consequences as a complete loss of presenilin function; 3) that loss of pen-1+ and pen-2+
20 function in a sel-12+ background results in phenotypes indicative of a partial loss of presenilin/Notch pathway function; 4) that pen-1 and pen-2 interact genetically with both sel-12 and hop-1; 5) that the open reading frames for pen-1 and pen-2 encode unrelated integral membrane proteins; 6) that pen-1 and pen-2 related genes are conserved across phyla.

Pen-1 and pen-2 mutations enhance sel-12 Δ to the lin-12/glp-1-like phenotypes
25 associated with total presenilin loss. As double mutants with a sel-12 Δ mutation, pen-1 alleles and pen-2 alleles each result in a set of phenotypes identical with those seen in sel-12 Δ ; hop-1 Δ worms that receive no maternal sel-12+ activity. Specifically, each of these double mutants with sel-12 Δ share 3 common abnormalities that are not seen in sel-12 Δ or hop-1 Δ single mutants, or in pen-1 or pen-2 single mutants. First, all three sets of double
30 mutants display indistinguishable glp-1-like sterile phenotypes characterized by germ cell proliferation defect similar to that described for glp-1 loss-of-function mutants (Austin and Kimble, 1987). Second, all three double mutants show a common cell fate specification defect(s) that indicates a loss of lin-12/Notch signaling. lin-12+ activity is required for the ventral uterine precursor versus anchor cell fate decision: lin-12(lf) mutants have 2 anchor
35 cells rather than the normal complement of one because the cell that normally adopts the

ventral uterine precursor fate instead becomes an anchor cell (Greenwald, I. et al., Cell (1983) 34:435-444). The sel-12 Δ ; pen-1 and sel-12 Δ ; pen-2 double mutants display this "2 anchor cell" phenotype just as does the sel-1 Δ 2; hop-1 Δ double mutant (Westlund, B et al., Proc. Natl. Acad. Sci. (1999) 96: 2497-2502). Third, sel-12 Δ ; pen-1 and sel-12 Δ ; pen-2 double mutants, like sel-12 Δ ; hop-1 Δ , display an everted vulva phenotype that is reminiscent of vulva defects seen in lin-12(lf) mutants.

The above phenotypic comparisons demonstrate that a reduction in pen-1+ or pen-2+ activity, in combination with a loss of sel-12+ activity, results in a loss of presenilin pathway function comparable to the effects of eliminating the two redundant presenilins encoded by sel-12 and hop-1.

Single pen-1 and pen-2 mutations confer phenotypes associated with partial loss of presenilin function. As single mutants, pen-1 and pen-2 worms display two visible abnormalities. First, pen-1 and pen-2 homozygotes (produced from a pen-1/+ or pen-2/+ mother) both produce normal numbers of self-progeny embryos but these embryos are retained in the animal's uterus and never laid. pen-1 and pen-2 hermaphrodites are thus egg-laying defective (or Egl), a phenotype shared by the sel-12 Δ single mutant.

Second, the embryos produced by homozygous pen-1 or pen-2 hermaphrodites never hatch but instead arrest in development with multiple abnormalities. The arrested embryos produced by pen-1 and pen-2 hermaphrodites show very similar abnormalities. Most strikingly, many of the arrested embryos make only a partial pharynx: the posterior pharyngeal lobe is present, but the anterior lobe is absent. Absence of anterior pharynx, called an Aph phenotype (for no anterior pharynx), was first described for certain weak alleles of glp-1. The GLP-1 receptor is required for a specific embryonic signaling event that induces formation of anterior pharynx (Mello, C. et al., Cell (1994) 77: 95-106; Moscovitz, I et al., Development (1994) 120:3325-3338; Hutter, H. and Schnabel, R., Development (1994) 120:2051-2064); absence of maternally provided glp-1+ activity can thus result in the Aph phenotype as well as other defects (Priess, J. et al., Cell (1987) 5:601-611). A connection of the Aph phenotype with reduced presenilin function comes from analysis of sel-12 Δ ; hop-1 Δ hermaphrodites which receive maternal sel-12+ (which rescues the sterility seen in the absence of maternal sel-12+ function). In this situation, sel-12 Δ ; hop-1 Δ hermaphrodites produce arrested embryos which display the Aph phenotype, as well addition to other glp-1like embryonic defects (Westlund, B., supra). These properties of pen-1 and pen-2 indicate both genes act in concert with both sel-12 and hop-1 presenilins, since the loss of pen-1 or pen-2 causes phenotypes more severe than those cause by the sel-12 or hop-1 single mutant.

In addition to the Aph phenotype, embryos produced by homozygous pen-1 or pen-2 hermaphrodites display other abnormalities. The embryos usually arrest with little evidence of elongation and the embryonic hypodermis (layer of epidermal cells that lies under and secretes the cuticle) often fails to fully enclose other cell types. Similar phenotypes have been described for embryos produced by glp-1(ts) mutants.

5 In summary, pen-1 and pen-2 mutants share multiple phenotypes (Egl, Aph and defective embryonic elongation) that are indicative of cell signaling defects involving the Notch family receptors glp-1 and lin-12. In addition, in combination with sel-12Δ, pen-1 and pen-2 result in additional, stronger Notch pathway-related defects (glp-1-like sterility, 2 anchor cell phenotype; vulva eversion). The combined genetic and phenotypic evidence indicates that pen-1 and pen-2 are novel components that may assist presenilins in Notch
10 receptor maturation and/or processing.

Pen-1 corresponds to the predicted *C. elegans* gene VF36H2L.1. To clone pen-1, we genetically mapped pen-1(ep140) to increasingly smaller intervals, first using visible genetic markers and then using molecular markers [Tc1 transposon insertions and single nucleotide polymorphisms (SNPs)]. The final stage of SNP mapping of pen-1 narrowed its position to a
15 52 KB interval on chromosome I. This interval, as documented in the *C. elegans* database ACEDB (Eeckman, F. and Durbin, R. *C. elegans: Modern Biological Analysis of an Organism* (1995) pp. 583-599), contains a total of 7 predicted genes. One of these, VF36H2L.1, was identified as pen-1 on the basis of RNA-mediated interference (RNAi) data and mutation detection. For many *C. elegans* genes, RNAi disrupts both maternal and
20 zygotic gene activity (Tabara, H. et al. *Science* (1998) 282:430-431). In case of pen-1, disruption of maternal activity after injection of dsRNA into adult hermaphrodites was evidenced by the production of developmentally arrested embryos with an Aph phenotype. As expected, this phenotype was observed after RNAi of both wild-type and sel-12Δ
25 hermaphrodites. RNAi in either background also gave many viable escaper progeny that grew to adulthood. In the case of RNAi in a sel-12Δ background, a high proportion of these escapers displayed glp-1-like sterility, consistent with inhibition of zygotic pen-1 activity. Unexpectedly, RNAi of VF36H2L.1 in wild type also resulted in Glp sterile progeny, although at a much lower frequency than in with pen-1 RNAi in a sel-12Δ homozygotes. By
30 contrast, glp-1-like sterility is never observed in pen-1 single mutants. This difference is most likely attributable to the property that RNAi typically disrupts both maternal and zygotic gene function, and can therefore result in more severe phenotypes than seen in zygotically lethal mutations.

By sequence analysis, we determined that the four pen-1 alleles isolated as sel-12
35 enhancers each contain single-nucleotide substitutions in the VF36H2L.1 open reading frame.

Remarkably, these 4 independently-derived lesions are each nonsense mutations in the same codon, Trp191. Three alleles (ep140, ep168, and ep170) are third base UGG to UGA alterations, while the fourth (ep216) is a second base UAG to UGG change. That these lesions result in Aph and glp-1 like sterility phenotypes similar to RNAi of VF36H2L.1, indicates that they are reduction-of- function mutations.

5 Pen-1 encodes an evolutionarily conserved protein with multiple transmembrane domains. The pen-1 (VF36H2L.1, GI#2815036) open reading frame is split among 4 exons that, when spliced, encode a 308 amino-acid protein. We confirmed the predicted splice junctions of exons 2 and 3 by sequence analysis of partial cDNA product. Pen-1 shows
10 homology with the predicted structures of various human, mouse, and Drosophila proteins, as described in detail below. Pen-1 is a predicted integral membrane protein that, as determined by the structure predicting programs PSORT2 and TopPred2, may contain up to 7 membrane-spanning domains.

Pen-2 corresponds to predicted *C. elegans* gene T28D6.9. We genetically mapped pen-2 to chromosome III between the cloned genes pha-1 and dpy-18. This interval spans
15 approximately 240 KB of DNA and contains 31 predicted genes as documented in ACEDB ver. 9 (Eeckman, F. and Durbin, R., *C. elegans: Modern Biological Analysis of an Organism* (1995) pp. 583-599). pen-2 was identified as the predicted gene T28D6.9 on the basis of RNAi data and mutation detection. RNAi of most of the genes in the interval led to the identification of, T28D6.9, as the only candidate gene for which RNAi gave the expected
20 maternal and zygotic pen-2 phenotypes. Wild type and sel-12Δ hermaphrodites injected with T28D6.9 produced a high proportion of developmentally-arrested embryos, many of which were Aph. In addition, RNAi of sel-12Δ (but not wild type) worms resulted in viable "escaper" progeny that displayed glp-1 like sterility. Mutation detection for the three pen-2 alleles isolated as sel-12 enhancers revealed that each contains a nonsense mutation in the
25 T28D6.9 predicted open reading frame. Two lesions (ep219 and ep220) alter the Trp74 codon, changing it from UGG to UGA (ep219) or UAG (ep220), while the third lesion (ep221) changes Trp36 to a UGA stop codon. These nonsense alleles should strongly reduce or abolish gene function, indicating that enhancement of sel-12Δ results from a loss of wild-type pen-2⁺ activity.

30 Pen-2 encodes a predicted multi-pass membrane protein. The pen-2 (T28D6.9, GI#3873415) open reading frame encodes a 101 amino acid protein. The predicted exon/intron structure of pen-2 has been confirmed by the sequence of an unpublished full-length cDNA (yk569h5 GI# 5572325 and 5558557) present in Genbank. Pen-2 shows a high level of homology with the predicted structures of various human, mouse, rat, and
35 Drosophila proteins, as described in detail below. Pen-2 is a predicted integral membrane

protein that, as determined by the structure predicting programs PSORT2 and TopPred2, contains 2 likely transmembrane domains.

Based on several properties, including their own specific phenotypes and their interactions with sel-12, pen-1 and pen-2 likely encode products that interact with presenilins. By extension, other genes with properties in common with pen-1, pen-2, sel-12, or hop-1 can be considered as potential presenilin interacting genes. We have identified the aph-2 gene as a presenilin interacting gene based on 1) the specific phenotypes associated with a loss of aph-2+ function and 2) our identification of novel genetic interactions of aph-2 and with sel-12 and hop-1, and with pen-1 and pen-2.

The aph-2 gene was identified by C. Goutte et al. (1995 International Worm Meeting, abstract 39; 1998 East Coast Worm Meeting, abstract 151; Worm Breeder's Gazette 12(5):27 (1993); Worm Breeder Gazette 13(d):83 (1994)) as a possible component of glp-1 mediated signaling in *C. elegans* embryos. The aph-2 mutants characterized by these investigators have no reported zygotic phenotypes, but do have maternal embryonic defects, including an Aph phenotype, strikingly similar to glp-1(ts) embryonic defects. aph-2 reportedly corresponds to the predicted gene ZC434.6. The predicted aph-2 protein is a 721 amino acids in length and is characterized by a signal sequence and 1 to 3 transmembrane domains as predicted by PSORT2 (Nakai K., and Horton P., Trends Biochem Sci, 1999, 24:34-6) and TopPred2 (Claros MG, and von Heijne G. Comput Appl Biosci 1994 Dec;10(6):685-6).

The screen that identified pen-1 and pen-2 did not yield mutations in aph-2. In order to identify potential presenilin/ aph-2 interactions that may have been missed due to the high stringency of this screen, we investigated a variety of genetic backgrounds that are more highly sensitized for presenilin loss. For these experiments, due to the lack of an available aph-2 mutations, we used RNAi to reduce aph-2+ function in selected backgrounds. Injection of aph-2 dsRNA into the germ line of wild type hermaphrodites results in highly penetrant embryonic lethality among the progeny, with many of the arrested embryos displaying an Aph phenotype. However, injected hermaphrodites still produce a substantial fraction of viable progeny that grow to adulthood with no phenotypic abnormalities in somatic tissues. These worms can be considered "transient escapers" because many of them produce developmentally-arrested Aph embryos. We were thus able to inject adult hermaphrodites with aph-2 RNA and examine their transient escaper progeny for presenilin-dependent phenotypes. Table 4 summarizes the results of these experiments.

Table 4 Enhancement of presenilin and pen gene phenotypes by aph-2 RNAi (transgene escaper progeny phenotypes).

Genotype1	Zygotic Phenotypes (no aph-2 RNAi)	Zygotic phenotypes after aph-2 RNAi 2
wild type	wild-type	wild-type
sel-12Δ	Egg-laying defective	Egg-laying defective
sel-12Δ; hop-1(ep168)/+	Egg-laying defective	glp-1 like sterility (12%)
hop-1(ep171)3	Low penetrance glp-1 like sterility (<1%)	glp-1 like sterility (>50%)
pen-1(ep140)	Egg-laying defective	glp-1 like sterility (>50%)
pen-2(ep220)	Egg-laying defective	glp-1 like sterility (>50%)

1 Complete genotypes of XX hermaphrodites were as follows: Row 1: N2 (wild-type). Row 2:

10 sel-12(ep6). Row 3: sel-12(ep6); hop-1(ep168) unc-74(x19)/ hT2 [hop-1+ unc-74+]. Row 4: hop-1(ep171) unc-74(x19). Row 5: unc-29(e1072) pen-1(ep140) Row 6: pen-2(ep220) dpy-18(e364).

15 2 Hermaphrodites from homozygous or heterozygous stocks were injected with aph-2 dsRNA.

All genotypes segregated developmentally-arrested Aph-2 embryos, as well as some viable escaper progeny. For genotypes where aph-2(RNAi) resulted in a zygotic Glp-1-like sterility among the escaper progeny, the approximate fraction of Glp sterile worms is indicated.

20 Aph-2 RNAi in a homozygous sel-12Δ background does not obviously enhance sel-12 to more severe presenilin phenotypes. However, significant enhancement is detected in homozygous sel-12Δ worms that are also heterozygous for a hop-1 nonsense mutant (hop-1(ep168) /+). With aph-2 RNAi, about 12% of sel-12Δ; hop-1(ep168)/ + animals display glp-1-like sterility, something never seen for this genotype in the absence of aph-2
25 RNAi. Further, the aph-2 RNAi enhancement is dependent on combined reduction in both presenilins as no interaction is seen with the sel-12Δ.

An additional interaction with aph-2 is observed with an unusual hop-1 allele, ep171. This allele carries a D to N missense alteration in an conserved aspartate residue that

corresponds to the Asp385 residue (located in TM domain 8) of human PS1. A PS1 Asp285Ala mutation results in loss of PS1 function, and also has dominant negative effects on PS1+ expression (Wolfe, M. Nature (1999) 398: 513-517). Like the PS1 D385A alteration, hop-1(ep171) has no wild type presenilin activity: the sel-12 Δ ; hop-1(ep171) double mutant has a glp-1-like sterility defect similar to sel-12 Δ ; hop-1 Δ . In a sel-12+ background, hop-1(ep171) results in a very low penetrance (<1%) glp-1-like sterility phenotype, which suggests it must have dominant negative effects on sel-12 presenilin function or expression. We found that RNAi of aph-2 in homozygous hop-1(ep171) hermaphrodites results in highly penetrant glp-1-like sterility (>50% of viable progeny), indicating a strong additive interaction between reduced aph-2+ function and the hop-1(ep171) dominant effects.

Finally, we also observed that aph-2 RNAi strongly enhances pen-1 and pen-2 mutant phenotypes. Homozygous adult pen-1 and pen-2 hermaphrodites segregating from heterozygous stocks have a normal-appearing germline and never exhibit glp-1-like sterility. In contrast, the corresponding pen-1; aph-2(RNAi) and pen-2; aph-2(RNAi) hermaphrodites display glp-1-like sterility at high penetrance (>50% of viable pen-1 or pen-2 homozygous progeny). These observations demonstrate that a variety genetic backgrounds with partially reduce presenilin pathway activity can be enhanced to stronger phenotypes by an RNAi-mediated reduction in aph-2 activity. The data demonstrate a functional interaction of aph-2 with presenilins and pen-1 and pen-2.

Structure of APH-2 and APH-2 related human and fly proteins. APH-2 contains a PSORT2 predicted cleavable signal sequence and 1 to 3 transmembrane domains predicted by PSORT2 and TopPred2. APH-2 is 18% identical in amino acid sequence to the predicted human protein encoded by the nearly full-length cDNA KIAA0253 (Nagase, T. et al. DNA research (1996) 3: 321-329). In addition, APH-2 shows a similar level of identity to a Drosophila protein predicted from contigged ESTs generated at Exelixis, Inc. The human and Drosophila APH-2 related proteins are 30% identical and Clustal alignments of the 3 proteins show conservation over entire length of each protein.

Methods: RNA mediated interference (RNAi). RNAi of specific genes was generally done using dsRNA prepared from templates of PCR-amplified genomic DNA fragments. The 5' end of the PCR primers contained the promoter sequences for T7 RNA polymerase and the 3' regions were designed such that they amplified one or more exons of the targeted gene. PCR reactions, employing 5mmole of each primer, and 0.5 mg of wild-type genomic in a 50ml reaction, were done using the Expand kit (Roche Biochemicals, Summerville, NJ), according to the manufacture's protocols. The PCR conditions were as follows: an initial denaturation at 95 C for 30sec, followed by 35 cycles of 94 C for 30sec, 55 C for 15 sec, 72 C for 1min, and a final extension at 72 C for 3min. Amplified DNA was ethanol precipitated

and resuspended in 20ml of RNase-free water. A portion of the PCR product was used as template for a T7 polymerase-directed in vitro transcription reaction according to the manufacturer's instructions (Promega, Inc). Reactions were precipitated with ethanol and RNA was resuspended in 20 ml of RNase-free water and 10 ml 3X IM buffer (20mM KPO₄ pH7.5, 3mM K+Citrate pH 7.5, 2% PEG 6000). The complementary sense- and anti- sense RNAs were annealed by incubation at 68 C for 10 minutes, followed by incubation 37 C for 30 minutes, and then centrifuged through a 0.45 um cellulose acetate filter. Microinjection of RNA was done as described (Fire et al., Development (1991) 113:503-514) using hermaphrodites at the L4 or young adult stage. Injected worms were recovered in M9 buffer (per liter: 30g Na₂HPO₄, 15g KH₂PO₄, 2.5g NaCl, 5g NH₄Cl) for 10-30 minutes, transferred to individual plates, and then transferred to new plates daily. The first generation self-progeny of injected hermaphrodites were inspected for RNAi induced phenotypes by observation in the dissecting microscope or in the compound microscope equipped with Nomarski differential interference optics.

C. elegans strains used. Methods for handling and culturing *C. elegans* have been described (Brenner, S. Genetics (1974) 77: 71-94). *C. elegans* variation. Bristol strain N2 represents wild type and is largely isogenic with most of the mutant strains used here. Specific mutations used for genetic mapping and characterization included: LG I - unc-74(x19), dpy-5(e61), unc-29(e1072), fog-3(q443), dpy-24(s71). LG III - dpy-19(e1259ts), unc-119(e2498), pha-1(e2123), dpy-18(n499 or e364). LG IV - him-8(e1489). LG X lon-2(e678). Rearrangements: mnDp66 (X; I). All are described in *C. elegans* II. Deletion mutations that remove most or part of the of sel-12 or hop-1 coding region are described below. Because the sel-12 gene is sex-linked and sel-12 mutants are mating defective, the transfer of sel-12Δ between strains was usually accomplished using males that carry the chromosomal duplication mnDp66 (X; I) which carries a complementing sel-12+ allele.

SNP Screening by DHPLC: Candidate SNPs were amplified separately from CB4856 and N2 genomic DNA. The PCR products were mixed, denatured and reannealed to create heterozygote molecules for screening by Denaturing HPLC (DHPLC). Each SNP was screened at 5 different temperatures using the same separation gradient. A SNP was deemed authentic when a heteroduplex was detected in the heterozygous state but not in the homozygous starting strains. The appropriate temperature for each SNP was noted and used for screening that SNP on recombinant worms.

SNP Scoring on recombinant worms: Lysates from appropriate recombinants were used as genomic DNA templates for amplifying the SNPs by PCR. These crude PCR products were then run on DHPLC using the appropriate temperatures for each SNP identified above.

For each recombinant, each SNP was typed and the data input into a spreadsheet at random. The physical order of the SNPs was then determined from AceDB. This generated a haplotype for each recombinant, and the locations at which recombination events occurred was noted.

Isolation and characterization of sel-12 and hop-1 deletion mutant strains. Deletion alleles of sel-12 and hop-1 were obtained by the two-step method of Plasterk (Plasterk, R. C. elegans: Modern Biological Analysis of an Organism (1995) pp. 59-80) using mut-2 as a source of Tc1 transposon mutator activity. sel-12(ep6) (hereinafter referred to as sel-12Δ) is a deletion mutation that removes amino acids 34 to 441 of the sel-12 open reading frame. hop-1(ep90) (hereinafter referred to as hop-1) is a 722 bp deletion that starts at amino acid 216 in the hop-1 open reading frame and terminates within the gene's 3' untranslated region. The sel-12Δ and hop-1Δ single mutants were backcrossed at least 10 times to wild-type (C. elegans variation. bristol strain N2, supra) before phenotypic characterization and construction of double mutants. The sel-12Δ single mutant has an egg-laying defective phenotype similar to that of previously-described sel-12(lf) mutants (Levitan, D. and Greenwald, I. Nature (1995) 377:351-354). The hop-1Δ mutant has no gross phenotypic abnormalities, similarly to hop-1 deletion alleles described by others (Westlund, B. et al. Proc. Natl. Acad. Sci (1999) 96: 2497-2502).

To provide a source of sel-12Δ; hop-1Δ double mutants that lack maternal sel-12+ activity, we constructed a balanced sel-12Δ/sel-12Δ; hop-1Δ +/+ unc-74 strain. This strain segregates doubly mutant sel-12Δ; hop-1Δ hermaphrodites that exhibit a completely penetrant sterile phenotype with germline proliferation defects characteristic of glp-1(lf) mutants (Austin, J. and Kimble, J., Cell (1987) 51:589-599). In addition, these worms have a fully penetrant 2 anchor cell phenotype and an everted vulva phenotype that is reminiscent of vulval defects caused by lin-12(lf) mutations.

Isolation of enhancers of sel-12Δ. Enhancer alleles of pen-1 and pen-2 were obtained after mutagenesis of a homozygous sel-12Δ strain or, in later experiments, a sel-12Δ; unc-74(x19) strain (the unc-74 mutation lies near hop-1 on chromosome I and was included to provide a built-in mapping resource). XX hermaphrodites of either genotype were mutagenized with ethyl methane sulfonate as described (Brenner, S. Genetics (1974) 77: 71-94). In the F1 generation, one (or sometimes two) hermaphrodites were picked onto individual growth plates (approximately 55,000 plates total). Three to five days later, the plates were screened for the appearance of sterile F2 progeny with a "dark" appearance indicative of a defect in germline proliferation. Candidate sterile mutants were then screened by Nomarski difference interference microscopy to identify those which exhibit glp-1 like sterility similar to sel-12Δ; hop-1Δ worms.

A set of 44 candidates identified in this way were subjected to a cross scheme designed to determine whether or not the sterile phenotype in these mutants was dependent on the worm's sel-12 genotype as would be expected for a sel-12 enhancer. For this test, each candidate was crossed to dpy-19 III; him-8; lon-2 males and the resulting cross-progeny were picked onto individual plates. In the following generation, the presence of sterile lon-2/lon-2 progeny (which are sel-12+/sel-12+ in the absence of recombination) indicate that sterile phenotype was not dependent on a loss of sel-12+ activity and was possibly due to a mutation in one of the known glp-1 pathway genes (glp-1, lag-1, lag-2). 29 candidates analyzed in this way were sel-12 independent and thus were rejected as possible presenilin enhancers. For 26 of the 29 rejected candidates, the mutation causing sterility segregated in trans to dpy-19, which is the expected behavior for a glp-1 allele. 9 of these LG III mutations failed to complement the sterile phenotype of known glp-1 alleles; the other 17 LG III mutations were not tested.

For the remaining 15 candidates, the Glp sterile phenotype did not reappear in the F2 generation, a result consistent with the presence of an enhancer mutation whose interaction with sel-12 is rescued by maternal sel-12+ activity. This explanation was tested by picking sel-12/sel-12 worms in the F2 generation onto individual plates and examining their progeny for reappearance of glp-1-like sterility in the next generation. This was the result observed for each of the remaining 15 candidates. A combination of complementation tests, meiotic mapping, and sequence analysis of mutant alleles demonstrated that each candidate carried a mutation in either hop-1 (8 candidates) or in either of two newly-identified genes, pen-1 (4 candidates) or pen-2 (3 candidates).

Pen-1 mapping, characterization, cloning, and computational analysis: Genetic mapping of pen-1 was done in sel-12Δ backgrounds and was based on the glp-1-like sterility phenotype of doubly mutant pen-1; sel-12Δ worms. We initially mapped pen-1(ep140) to chromosome I between unc-29 and dpy-24. Further mapping with visible markers narrowed the position to between unc-29 and fog-3, a 1.1 MB interval. From heterozygotes of the genotype pen-1/unc-29 fog-3 trans-heterozygotes, 16/20 Unc-29 non-Fog-3 recombinants and 1/4 Fog-3 non-Unc-29 recombinants segregated pen-1.

Finer mapping was done using SNP markers that are polymorphic between the N2 Bristol strain from which pen-1 mutants were derived and strain CB4856 Hawaiian strain of *C. elegans*. The Genome Sequencing Center (St. Louis, MO) has identified a large number of potential SNPs in CB4856 (<http://genome.wustl.edu/gsc/CEpolymorph/snp.shtml>). Four of these potential SNPs in the unc-29 to fog-3 interval were confirmed by testing with an SNP genotyping assay that is based on separation of heteroduplex PCR products by denaturing HPLC (Underhill PA, et al., Genome Res. 1997 Oct;7(10):996-1005). Initial

mapping against these SNPs was done by constructing heterozygotes of the genotypes unc-29 pen-1/ CB4856 or pen-1 dpy-24/CB4956 and picking Unc-29 non-Pen-1 or Dpy-24 non-Pen-1 recombinants. Additional non-Pen-1 recombinants were isolated from unc-29 pen-1 fog-3/ CB4856 heterozygotes. Among 50 Unc-29 non-Pen-1 recombinants, 9 had cross-overs occurring to the right of the C31H5 SNP, placing pen-1 to the right of this marker. Among a combined set of 45 Dpy-24 non-Pen-1 or Fog-3 non-Pen-1 recombinants, 4 had cross-overs to the left of the F14B4 SNP, placing pen-1 left of this marker. The combined data positioned pen-1 to the ~240 KB interval between the C31H5 and F14B4 SNPs.

For mapping pen-1 to smaller intervals, additional SNPs lying in the C45G3 to F14B4 interval were identified by DNA sequencing. Six approximately 2 KB segments of DNA in the interval were amplified by PCR of N2 and CB4856 genomic DNA and end sequenced. In some cases, additional sequencing primers were used to generate internal sequence. SNPs between N2 and CB4856 were identified by sequence alignment and ~200 bp PCR products were designed around each high quality candidate. These 200 bp products were screened and scored by DHPLC as described above. This analysis positioned pen-1 between 2 SNPs, one on cosmid C45G3 and the other on cosmid F36H2, that lie approximately 52 KB from one another.

Mutation detection. Two single nucleotide polymorphisms (SNPs), labeled C45G3A and F36H2A, defined a 52kb genomic interval, pen-1, within which seven predicted candidate genes resided. A 30kb gene-rich section of this 52kb interval was resequenced in 3 worms whose mutation had been mapped genetically to this region, ep140, ep169, and ep170. All DNA sequencing reactions were performed using standard protocols for the BigDye sequencing reagents (Applied Biosystems, Inc. Foster City, CA) and products were analyzed using ABI 377 DNA sequencers.

Trace data obtained from the ABI 377 DNA sequencers was analyzed and assembled into contigs using the Phred-Phrap programs (Gordon, Genome Res. (1998) 8:195-202). The resequence data was then compared to the wildtype strain, N2, for polymorphism. This analysis identified a third position base change, G to an A, at 191AA in the VF36H2L.1 gene (GI# 2815036) in three pen-1 alleles, ep140, ep169 and ep170, resulting in an amino acid change from a tryptophan (W) to stop (*). Further sequencing analysis of unmapped mutants revealed another mutation in worm ep216 within the same codon, but in the second position, also a G to an A, resulting in the same amino acid change. Analysis of human pen-1 led us to identify the novel pen-1B protein.

Pen-2 mapping, characterization, cloning, and computational analysis. We initially positioned pen-2(ep220) to the left of unc-25 on chromosome III. From hermaphrodites of

the genotype *pen-2/ dpy-18 unc-25*, 4/4 non-Dpy-18 Unc-25 recombinants segregated *pen-2* and 0/21 Dpy-18 non-Unc-25 recombinants segregated *pen-2*. Of 70 *dpy-18 unc-25* homozygotes picked from the same heterozygous hermaphrodites, only 2 segregated *pen-2*, indicating that *pen-2* lies relatively close to *dpy-18* and probably to the left of this gene. Further mapping positioned *pen-1* between *pha-1* and *dpy-18*: from *pen-1/ pha-1 dpy-19; sel-12Δ* hermaphrodites, 1/14 non-Pha-1 Dpy-1 recombinants picked up *pen-2*. These data positioned *pen-2* between *pha-1* and *dpy-18* interval, an approximately 240 KB interval, and suggested *pen-2* lies close to *dpy-18*.

Identification of *pen-2* as predicted gene T28D6.9: We determined that *pen-2* is the predicted gene T28D6.9 (GI#3873415) based on 1) RNAi of predicted genes in the *pha-1* to *dpy-18* interval and 2) mutation detection. For 28 of the 31 genes in the interval, primers tailed with T7 promoter sequence were used to amplify selected coding region using either a first-strand cDNA pool or genomic DNA as template as described above. Double-stranded RNA was synthesized from each PCR product and injected into *sel-12Δ* homozygotes. RNAi of T28D6.9 produced the expected *pen-2* phenotypes among the progeny of injected worms, including *glp-1*-like sterility in *sel-12Δ* worms and an Aph embryonic arrest phenotype after injection into N2 and *sel-12Δ* worms. Mutation detection of the T28D6.9 open reading frame identified nonsense mutations in each of three *pen-2* alleles (*ep219*, *ep220*, *ep221*). Briefly, The single mutant in this group, *ep220*, was tested by sequencing a PCR product amplified in an *ep220* lysate and wild-type strain. This analysis identified a G to A mutation at 74AA that resulted in a tryptophan (W) to stop (*). Additional sequencing analysis of unmapped mutants revealed that there were 2 more mutants in this group. The *ep219* worm had a G to A change in the third position of 74AA that produced a W to a *. The *ep221* worm had a G to A change in another W that also resulted in a stop codon at 36AA. These three changes all effect highly conserved tryptophans that could significantly alter or ablate the function of the T28D6.9 gene.

V. Cell-based reporter assays.

We developed a cell culture gamma secretase assay based on a reporter construct carrying the C-terminal 99 amino acids of APP fused to a Gal4VP16 transcriptional activator protein. The Gal4 moiety is retained at the cell surface by the APP transmembrane domain until presenilin-dependent cleavage releases it to translocate to the nucleus and activate transcription of a UAS-luciferase reporter transgene. In assay validation experiments, a known gamma secretase inhibitor completely blocked reporter gene activity, and known dominant negative presenilin mutations also inhibited the reporter activity. A conceptually similar assay has been shown previously to work in *Drosophila* in vivo using a UAS-beta-galactosidase transgene reporter. Beta-galactosidase reporter gene activity, and hence

gamma-secretase-like protease activity have been shown to be absolutely dependent on the presence of presenilin in this in vivo assay.

Our data show that inhibition of pen-1, pen-2, aph-2 or presenilin results in a strong reduction of steady state presenilin protein levels, and a correlated reduction of gamma-secretase activity. The effects of pen-1 or aph-2 inhibition is equally strong, and pen-2 nearly as strong, with respect to both reduction of gamma secretase activity and presenilin protein reduction as is presenilin inhibition itself. Presenilin reduction is verified in multiple cell-based systems, including Drosophila and human cell systems, using several inhibitors, including RNAi, heterocyclic compounds identified in the disclosed screens, and intrabodies. These data provide assays for a functional interaction of the pen genes and presenilin. Hence, the invention provides a method for specifically detecting a stress that alters a functional interaction of a presenilin enhancer (pen) polypeptide with a presenilin by introducing a predetermined stress into a system which provides a functional interaction of a pen polypeptide with a presenilin, whereby the system provides a stress-biased interaction of the pen polypeptide with the presenilin, wherein the absence of the stress, the system provides an unbiased interaction of the pen polypeptide with the presenilin; and detecting the stress-biased interaction of the pen polypeptide with the presenilin as a change in an amount of presenilin in the system, wherein the amount may be expressed as an amount of presenilin N or C-terminal fragments (NTFs), presenilin holoprotein, or a ratio thereof, wherein a difference between the stress-biased and unbiased interactions indicates that the stress alters the interaction of the pen polypeptide with the presenilin.

All publications and patent applications cited in this specification are herein incorporated by reference as if each individual publication or patent application were specifically and individually indicated to be incorporated by reference. Although the foregoing invention has been described in some detail by way of illustration and example for purposes of clarity of understanding, it will be readily apparent to those of ordinary skill in the art in light of the teachings of this invention that certain changes and modifications may be made thereto without departing from the spirit or scope of the appended claims.

WHAT IS CLAIMED IS:

1. A method for specifically detecting a stress that alters a functional interaction of a presenilin enhancer (pen) polypeptide with Notch or APP processing, the method comprising steps:

5 introducing a predetermined stress into a system which provides a functional interaction of a pen polypeptide with Notch or APP processing, whereby the system provides a stress-biased interaction of the pen polypeptide with Notch or APP processing, wherein the absence of the stress, the system provides unbiased interaction of the pen polypeptide with Notch or APP processing; and

10 detecting the stress-biased interaction of the pen polypeptide with Notch or APP processing, wherein a difference between the stress-biased and unbiased interactions indicates that the stress alters the interaction of the pen polypeptide with Notch or APP processing;

15 wherein the system is selected from the group consisting of: (i) a viable cell expressing the pen polypeptide wherein the pen polypeptide expression is determined to be non-natural or pathogenic, and (ii) an in vitro, cell-free mixture comprising a determined amount of the pen polypeptide,

20 wherein the pen polypeptide has sequence similarity to a pen polypeptide selected from the group consisting of: human, rat, mouse, D. melanogaster and C. elegans pen-2; human, mouse, D. melanogaster and C. elegans pen-1; and human pen-1B, wherein the similarity is at least 20% identity.

2. A method for specifically detecting a stress that alters a functional interaction of an enhancer of presenilin (pen) polypeptide with APP processing, the method comprising steps:

25 introducing a predetermined stress into a system which provides a functional interaction of a pen polypeptide with APP processing, whereby the system provides a stress-biased interaction of the pen polypeptide with APP processing, wherein the absence of the stress, the system provides unbiased interaction of the pen polypeptide with APP processing; and

30 detecting the stress-biased interaction of the pen polypeptide with APP processing, wherein a difference between the stress-biased and unbiased interactions indicates that the stress alters the interaction of the pen polypeptide with APP processing;

wherein the system is selected from the group consisting of: (i) a viable cell expressing the pen polypeptide wherein the pen polypeptide expression is determined to be

non-natural or pathogenic, and (ii) an in vitro, cell-free mixture comprising a determined amount of the pen polypeptide,

wherein the pen polypeptide has sequence similarity to a pen polypeptide selected from the group consisting of: human, rat, mouse, *D. melanogaster* or *C. elegans* pen-2; human, mouse, *D. melanogaster* or *C. elegans* pen-1; human pen-1B; and human, *D. melanogaster* or *C. elegans* Aph-2, wherein the similarity is at least 20% identity.

3. A method for specifically detecting a stress that alters a functional interaction of a presenilin enhancer (pen) polypeptide with a presenilin, the method comprising steps:

introducing a predetermined stress into a system which provides a functional interaction of a pen polypeptide with a presenilin, whereby the system provides a stress-biased interaction of the pen polypeptide with the presenilin, wherein the absence of the stress, the system provides an unbiased interaction of the pen polypeptide with the presenilin; and

detecting the stress-biased interaction of the pen polypeptide with the presenilin as a change in an amount of presenilin in the system, wherein the amount may be expressed as an amount of presenilin N or C-terminal fragments (NTFs), presenilin holoprotein, or a ratio thereof, wherein a difference between the stress-biased and unbiased interactions indicates that the stress alters the interaction of the pen polypeptide with the presenilin;

wherein the system is selected from the group consisting of: (i) a viable cell expressing the pen polypeptide wherein the pen polypeptide expression is determined to be non-natural or pathogenic, and (ii) an in vitro, cell-free mixture comprising a determined amount of the pen polypeptide,

wherein the pen polypeptide has sequence similarity to a pen polypeptide selected from the group consisting of: human, rat, mouse, *D. melanogaster* and *C. elegans* pen-2; human, mouse, *D. melanogaster* and *C. elegans* pen-1; and human pen-1B, wherein the similarity is at least 20% identity.

4. A method according to claim 1, 2 or 3, wherein the pen polypeptide is selected from the group consisting of: human, rat, mouse, *D. melanogaster* and *C. elegans* pen-2.

5. A method according to claim 1, 2 or 3, wherein the pen polypeptide is selected from the group consisting of: human, mouse, *D. melanogaster* and *C. elegans* pen-1.

6. A method according to claim 1, 2 or 3, wherein the pen polypeptide is human pen-1B.

7. A method according to claim 1, 2 or 3, wherein said the pen polypeptide is a naturally-occurring pen polypeptide identifiable in a sel-12 Δ (delta) homozygous *C. elegans* genetic mutation enhancer screen.

5

8. A method according to claim 1, 2, 3, 4, 5, 6 or 7, wherein the identity is at least 50%.

9. A method according to claim 1, 2, 3, 4, 5, 6 or 7, wherein the identity is 100%.

10

10. A method according to claim 1, 4, 5, 6, 7, 8 or 9, wherein the functional interaction comprises binding of the pen polypeptide with a component of Notch or APP processing.

15

11. A method according to claim 1, 4, 5, 6, 7, 8 or 9, wherein the functional interaction comprises binding of the pen polypeptide with a component of Notch or APP processing and the component is a presenilin or γ -secretase.

12. A method according to claim 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or 11 wherein the system is the viable cell and the stress is a pharmacologically active agent.

20

13. A method according to claim 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or 11 wherein the system is the viable cell and the stress is a deficiency in functional expression of the pen polypeptide.

25

14. A method according to claim 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or 11 wherein the system is the viable cell and the stress is a deficiency in functional expression of the pen polypeptide by virtue of genomic disruption of otherwise endogenous alleles encoding the pen polypeptide.

30

15. A method according to claim 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or 11 wherein the system is the viable cell and the stress is a deficiency in functional expression of the pen polypeptide by virtue of coexpression of a polynucleotide comprising a sequence antisense of an endogenous allele encoding the pen polypeptide.

16. A method according to claim 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or 11, wherein the system is the viable cell and the cell is in situ (resident in an animal host).

17. A method according to claim 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or 11, wherein the system is the viable cell and the cell is in vitro (isolated from an animal host).

5

18. A method according to claim 1, 2, 4, 5, 6, 7, 8, 9, 10, or 11, wherein the system is the in vitro, cell-free mixture and the stress is a pharmacologically active agent.

10

19. A method according to claim 1, 2, 4, 5, 6, 7, 8, 9, 10, or 11, wherein the detecting step detects an indication of Alzheimer's disease.

20. A method according to claim 1, 2, 4, 5, 6, 7, 8, 9, 10, or 11, wherein the detecting step detects a transcriptional reporter of notch.

15

21. A method according to claim 1, 2, 4, 5, 6, 7, 8, 9, 10, or 11, wherein the detecting step detects the generation of A β (beta).

22. A method according to claim 1, 2, 4, 5, 6, 7, 8, 9, 10, or 11, wherein the detecting step detects the generation of A β (beta) using an A β -specific antibody.

20

23. A method according to claim 1, 2, 4, 5, 6, 7, 8, 9, 10, or 11, wherein the detecting step detects a structural alteration in the pen polypeptide.

25

24. A method according to claim 1, 2, 4, 5, 6, 7, 8, 9, 10, or 11, wherein the detecting step detects a structural alteration in the pen polypeptide using a pen polypeptide-specific antibody.

25. A method according to claim 3, wherein the change in amount of detectable presenilin is detected by presenilin-specific antibodies.

30

26. A method according to claim 1, 2, 4, 5, 6, 7, 8, 9, 10, or 11,, wherein the system is the viable cell and comprises a gamma secretase reporter.

27. A method according to claim 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or 11, wherein the system is the viable cell and comprises a gamma secretase reporter comprising a C-terminal APP - Gal4 fusion protein and a UAS-reporter transgene, whereby cleavage of the fusion protein by gamma secretase releases Gal4, which in turn activates transcription of the transgene to express the reporter.

28. A method according to claim 2, 3, 4, 5, 6, 7, 8, 9, 10, or 11, wherein the pen polypeptide is selected from the group consisting of: human, D. melanogaster or C. elegans Aph -2.

29. An isolated polypeptide comprising a sequence having at least 50% sequence similarity to SEQ ID NO:1; wherein the polypeptide or fragment cross-reacts with a human pen-1B specific antibody.

30. A polypeptide according to claim 28 wherein the polypeptide binds a human presenilin or γ -secretase.

31. A polypeptide according to claim 29 or 30 comprising an amino acid sequence selected from the group consisting of:

SEQ ID NO:6, res 1-14;

SEQ ID NO:6, res 115-126;

SEQ ID NO:6, res 6-15;

SEQ ID NO:6, res 130-140;

SEQ ID NO:6, res 10-20;

SEQ ID NO:6, res 139-151;

SEQ ID NO:6, res 25-46;

SEQ ID NO:6, res 166-182;

SEQ ID NO:6, res 62-71;

SEQ ID NO:6, res 184-198;

SEQ ID NO:6, res 67-76;

SEQ ID NO:6, res 214-232; and

SEQ ID NO:6, res 72-95;

SEQ ID NO:6, res 246-257.

32. A polypeptide according to claim 29 comprising an amino acid sequence selected from the group consisting of: SEQ ID NOS:2-10.

33. A polypeptide according to claim 29 comprising an amino acid sequence selected from the group consisting of: SEQ ID NO:6, residues 1-254; SEQ ID NO:6, residues 4-255; SEQ ID NO:6, residues 9-257; and SEQ ID NO:6, residues 2-255.

34. A polypeptide according to claim 29 comprising SEQ ID NO:2.

35. A recombinant polynucleotide encoding a polypeptide according to claim 29, 30, 31, 32, 33 or 34..

36. A recombinant polynucleotide encoding a polypeptide according to claim 29, 30, 31, 32, 33 or 34 contained in a vector or a cell.

37. A method of making a polypeptide, said method comprising steps: introducing a polynucleotide according to claim 29, 30, 31, 32, 33 or 34 into a host cell or cellular extract, incubating said host cell or extract under conditions whereby said polynucleotide is expressed as a transcript and said transcript is expressed as a translation product comprising said polypeptide.

38. A method of screening for an agent that modulates the interaction of a pen-1B polypeptide to a binding target, said method comprising the steps of:

incubating a mixture comprising:

an isolated polypeptide according to claim 29, 30, 31, 32, 33 or 34,

a binding target of said polypeptide, and

a candidate agent;

under conditions whereby, but for the presence of said agent, said polypeptide specifically binds said binding target at a reference affinity;

detecting the binding affinity of said polypeptide to said binding target to determine an agent-biased affinity,

wherein a difference between the agent-biased affinity and the reference affinity indicates that said agent modulates the binding of said polypeptide to said binding target.

39. A method according to claim 38, wherein said binding target comprises a human presenilin or γ -secretase.

40. A method of screening for an agent that modulates the interaction of a pen-1B polypeptide to a binding target, said method comprising the steps of:

5 incubating a polynucleotide according to claim 29, 30, 31, 32, 33 or 34 under conditions whereby the polypeptide is expressed;

 incubating a mixture comprising said polypeptide, a binding target of said polypeptide, and a candidate agent under conditions whereby, but for the presence of said agent, said polypeptide specifically binds said binding target at a reference affinity;

10 detecting the binding affinity of said polypeptide to said binding target to determine an agent-biased affinity,

 wherein a difference between the agent-biased affinity and the reference affinity indicates that said agent modulates the binding of said polypeptide to said binding target.

15 41. A method according to claim 40, wherein said polynucleotide is in a cell.

 42. A method according to claim 40, wherein said binding target comprises a presenilin or γ -secretase.

SEQUENCE LISTING

<110> Curtis, Dan
 Francis, Ross
 5 McGrath, Garth Joseph
 Nicoll, Sharmon Monique
 Ruddy, David Andrew
 Ellis, Michael Christopher
 <120> Presenilin Enhancers
 10 <130> EX00-033
 <140>
 <141>
 <160> 38
 <170> PatentIn Ver. 2.1
 15
 <210> 1
 <211> 308
 <212> PRT
 <213> Caenorhabditis elegans
 20 <400> 1
 Met Gly Tyr Leu Leu Thr Ile Ala Cys Tyr Ile Ala Ser Phe Ser Pro
 1 5 10 15
 Ser Ile Ala Leu Phe Cys Ser Phe Ile Ala His Asp Pro Val Arg Ile
 20 25 30
 25 Ile Leu Phe Phe Leu Gly Ser Phe Phe Trp Leu Val Ser Leu Leu Phe
 35 40 45
 Ser Ser Leu Ala Trp Leu Gly Leu Ser Thr Val Leu Pro Asp Thr Phe
 50 55 60
 Leu Leu Ser Leu Thr Val Cys Ile Ile Ala Gln Glu Leu Ser Arg Val
 30 65 70 75 80
 Ala Tyr Phe Met Leu Leu Lys Lys Ala Gln Arg Gly Leu Asn Lys Ile
 85 90 95
 Thr Arg Gln Gly Gln Ile Ser Val Ala Pro Gly Val Ser Asp Leu His
 100 105 110
 35 Asn Ala Arg His Met Leu Ala Leu Val Cys Gly Leu Gly Met Gly Val
 115 120 125
 Ile Ser Ala Leu Phe Tyr Thr Met Asn Ala Phe Ala Ile Phe Ser Gly
 130 135 140
 Pro Gly Thr Ile Gly Leu Pro Asn Ala Leu Lys Thr Gly Glu Ile Asp

145 150 155 160
 Thr Asn Arg Ala Gly Lys Tyr Leu Pro Leu Cys Tyr Thr Leu Ser Ala
 165 170 175
 Ile Leu Leu Thr Leu Phe His Val Thr Trp Thr Ile Met Val Trp Asp
 180 185 190
 5 Ser Cys His Lys Ile Gly Arg Ile Pro Ser Ala Phe Val Pro Gly Ala
 195 200 205
 Ala Ala Val Val Ser His Leu Leu Val Thr Phe Leu Ser Ser Leu Asn
 210 215 220
 Ser Arg Gly Phe His Val Leu Val Phe Ala Val Gln Phe Leu Ile Leu
 10 225 230 235 240
 Leu Ile Cys Ile Ala Tyr Cys Asn Val Ile Met Gly Gly Thr Ile Ser
 245 250 255
 Ser Phe Val Asn Gly Ile Gly Gln Ser Ile Thr Asp Ala Val Thr Leu
 260 265 270
 15 Lys Gln Val Arg Thr Leu Ile Glu Glu Arg Lys Leu Arg Thr Gln Arg
 275 280 285
 Gln Ser Val Pro Asp Glu Pro Met Thr Glu Arg Ala Gly Thr Ser Asn
 290 295 300

 20 Thr Val Asn Ala
 305

 <210> 2
 <211> 238
 25 <212> PRT
 <213> Drosophila melanogaster
 <400> 2
 Met Thr Leu Pro Glu Phe Phe Gly Cys Thr Phe Ile Ala Phe Gly Pro
 1 5 10 15
 30 Pro Phe Ala Leu Phe Val Phe Thr Ile Ala Asn Asp Pro Val Arg Ile
 20 25 30
 Ile Ile Leu Ile Ala Ala Ala Phe Phe Trp Leu Leu Ser Leu Leu Ile
 35 40 45
 Ser Ser Leu Trp Tyr Ala Leu Ile Pro Leu Lys Glu Phe Leu Ala Phe
 35 50 55 60
 Gly Val Val Phe Ser Val Cys Phe Gln Glu Ala Phe Arg Tyr Ile Ile
 65 70 75 80
 Tyr Arg Ile Leu Arg Ser Thr Glu Gln Gly Leu His Ala Val Ala Glu

85 90 95
 Asp Thr Arg Val Thr Asp Asn Lys His Ile Leu Ala Tyr Val Ser Gly
 100 105 110
 Leu Gly Phe Gly Ile Ile Ser Gly Met Phe Ala Leu Val Asn Val Leu
 115 120 125
 5 Ala Asp Met Ser Gly Pro Gly Thr Met Gly Leu Lys Gly Gly Thr Glu
 130 135 140
 Leu Phe Phe Val Thr Ser Ala Ala Gln Ala Leu Ser Ile Ile Leu Leu
 145 150 155 160
 His Thr Phe Trp Ser Val Ile Phe Phe Asn Ala Phe Asp Thr Asn Asn
 10 165 170 175
 Tyr Ile His Ile Gly Tyr Val Val Phe Ser His Leu Phe Val Ser Leu
 180 185 190
 Ile Thr Leu Leu Asn Ala Asn Glu Leu Tyr Thr Thr Thr Leu Leu Ile
 195 200 205
 15 Asn Tyr Leu Val Thr Ile Leu Thr Gly Val Leu Ala Phe Arg Val Ala
 210 215 220
 Gly Gly Thr Ser Arg Ser Phe Arg Lys Phe Ile Thr Cys Gln
 225 230 235
 20 <210> 3
 <211> 126
 <212> PRT
 <213> *Heliothis virescens*
 <400> 3
 25 Met Thr Leu Ala Glu Phe Phe Ser Cys Ser Leu Leu Ala Phe Gly Ala
 1 5 10 15
 Pro Leu Val Met Phe Ala Leu Thr Val Ala Asn Asp Pro Val Arg Ile
 20 25 30
 Ile Ile Met Ile Ala Ala Ala Phe Gly Trp Leu Leu Ser Phe Leu Val
 30 35 40 45
 Ser Ser Val Val Trp Tyr Ala Val Val Pro Leu Arg Ser Tyr Leu Ala
 50 55 60
 Phe Gly Met Val Phe Ala Ile Ile Phe Gln Glu Val Phe Arg Tyr Gly
 65 70 75 80
 35 Met Tyr Val Leu Leu Arg Lys Thr Glu Ala Gly Leu Lys Glu Ile Ser
 85 90 95
 Glu Asn His Asn Ile Gly Ser Asn Lys Leu Glu Met Ala Tyr Val Ser

100 105 110
 Gly Leu Gly Phe Gly Thr Met Ser Gly Ala Phe Ala Leu Ile
 115 120 125

 <210> 4
 5 <211> 244
 <212> PRT
 <213> mouse
 <400> 4
 Met Gly Ala Ala Val Phe Phe Gly Cys Thr Phe Val Ala Phe Gly Pro
 10 1 5 10 15
 Ala Phe Ser Leu Phe Leu Ile Thr Val Ala Gly Asp Pro Leu Arg Val
 20 25 30
 Ile Ile Leu Val Ala Gly Ala Phe Phe Trp Leu Val Ser Leu Leu Leu
 35 40 45
 15 Ala Ser Val Val Trp Phe Ile Leu Val His Val Thr Asp Arg Ser Asp
 50 55 60
 Ala Arg Leu Gln Tyr Gly Leu Leu Ile Phe Gly Ala Ala Val Ser Val
 65 70 75 80
 Leu Leu Gln Glu Val Phe Arg Phe Ala Tyr Tyr Lys Leu Leu Lys Lys
 20 85 90 95
 Ala Asp Glu Gly Leu Ala Ser Leu Ser Glu Asp Gly Arg Ser Pro Ile
 100 105 110
 Ser Ile Arg Gln Met Ala Tyr Val Ser Gly Leu Ser Phe Gly Ile Ile
 115 120 125
 25 Ser Gly Val Phe Ser Val Ile Asn Ile Leu Ala Asp Ala Leu Gly Pro
 130 135 140
 Gly Val Val Gly Ile His Gly Asp Ser Pro Tyr Tyr Phe Leu Thr Ser
 145 150 155 160
 Ala Phe Leu Thr Ala Ala Ile Ile Leu Leu His Thr Phe Trp Gly Val
 30 165 170 175
 Val Phe Phe Asp Ala Cys Glu Arg Arg Arg Tyr Trp Ala Leu Gly Leu
 180 185 190
 Val Val Gly Ser His Leu Leu Thr Ser Gly Leu Thr Phe Leu Asn Pro
 195 200 205
 35 Trp Tyr Glu Ala Ser Leu Leu Pro Ile Tyr Ala Val Thr Val Ser Met
 210 215 220
 Gly Leu Trp Ala Phe Ile Thr Ala Gly Val Pro Ser Glu Val Ser Ser
 225 230 235 240

Ala Ala Phe Val

<210> 5

<211> 251

<212> PRT

5 <213> human

<400> 5

Met Gly Ala Ala Val Phe Phe Gly Cys Thr Phe Val Ala Phe Gly Pro
 1 5 10 15
 Ala Phe Ala Leu Phe Leu Ile Thr Val Ala Gly Asp Pro Leu Arg Val
 10 20 25 30
 Ile Ile Leu Val Ala Gly Ala Phe Phe Trp Leu Val Ser Leu Leu Leu
 35 40 45
 Ala Ser Val Val Trp Phe Ile Leu Val His Val Thr Asp Arg Ser Asp
 50 55 60
 15 Ala Arg Leu Gln Tyr Gly Leu Leu Ile Phe Gly Ala Ala Val Ser Val
 65 70 75 80
 Leu Leu Gln Glu Val Phe Arg Phe Ala Tyr Tyr Lys Leu Leu Lys Lys
 85 90 95
 Ala Asp Glu Gly Leu Ala Ser Leu Ser Glu Asp Gly Arg Ser Pro Ile
 100 105 110
 20 Ser Ile Arg Gln Met Ala Tyr Val Ser Gly Leu Ser Phe Gly Ile Ile
 115 120 125
 Ser Gly Val Phe Ser Val Ile Asn Ile Leu Ala Asp Ala Leu Gly Pro
 130 135 140
 25 Gly Val Val Gly Ile His Gly Asp Ser Pro Tyr Tyr Phe Leu Thr Ser
 145 150 155 160
 Ala Phe Leu Thr Ala Ala Ile Ile Leu Leu His Thr Phe Trp Gly Val
 165 170 175
 Val Phe Phe Asp Ala Cys Glu Arg Arg Arg Tyr Trp Ala Leu Gly Leu
 180 185 190
 30 Val Val Gly Ser His Leu Leu Thr Ser Gly Leu Thr Phe Leu Asn Pro
 195 200 205
 Trp Tyr Glu Ala Ser Leu Leu Pro Ile Tyr Ala Val Thr Val Ser Met
 210 215 220
 35 Gly Leu Trp Ala Phe Ile Thr Ala Gly Gly Ser Leu Arg Ser Ile Gln
 225 230 235 240
 Arg Ser Ser Cys Val Arg Thr Asp Tyr Leu Asp
 245 250

<210> 6

<211> 257

<212> PRT

<213> human

<400> 6

5 Met Thr Ala Ala Val Phe Phe Gly Cys Ala Phe Ile Ala Phe Gly Pro
 1 5 10 15
 Ala Leu Ala Leu Tyr Val Phe Thr Ile Ala Thr Glu Pro Leu Arg Ile
 20 25 30
 Ile Phe Leu Ile Ala Gly Ala Phe Phe Trp Leu Val Ser Leu Leu Ile
 10 35 40 45
 Ser Ser Leu Val Trp Phe Met Ala Arg Val Ile Ile Asp Asn Lys Asp
 50 55 60
 Gly Pro Thr Gln Lys Tyr Leu Leu Ile Phe Gly Ala Phe Val Ser Val
 65 70 75 80
 15 Tyr Ile Gln Glu Met Phe Arg Phe Ala Tyr Tyr Lys Leu Leu Lys Lys
 85 90 95
 Ala Ser Glu Gly Leu Lys Ser Ile Asn Pro Gly Glu Thr Ala Pro Ser
 100 105 110
 Met Arg Leu Leu Ala Tyr Val Ser Gly Leu Gly Phe Gly Ile Met Ser
 20 115 120 125
 Gly Val Phe Ser Phe Val Asn Thr Leu Ser Asp Ser Leu Gly Pro Gly
 130 135 140
 Thr Val Gly Ile His Gly Asp Ser Pro Gln Phe Phe Leu Tyr Ser Ala
 145 150 155 160
 25 Phe Met Thr Leu Val Ile Ile Leu Leu His Val Phe Trp Gly Ile Val
 165 170 175
 Phe Phe Asp Gly Cys Glu Lys Lys Lys Trp Gly Ile Leu Leu Ile Val
 180 185 190
 Leu Leu Thr His Leu Leu Val Ser Ala Gln Thr Phe Ile Ser Ser Tyr
 30 195 200 205
 Tyr Gly Ile Asn Leu Ala Ser Ala Phe Ile Ile Leu Val Leu Met Gly
 210 215 220
 Thr Trp Ala Phe Leu Ala Ala Gly Gly Ser Cys Arg Ser Leu Lys Leu
 225 230 235 240
 35 Cys Leu Leu Cys Gln Asp Lys Asn Phe Leu Leu Tyr Asn Gln Arg Ser
 245 250 255
 Arg

<210> 7

<211> 101

<212> PRT

<213> *Caenorhabditis elegans*

<400> 7

5 Met Asp Ile Ser Lys Leu Thr Asp Val Lys Lys Val Asp Leu Cys Lys
 1 5 10 15
 Lys Tyr Phe Leu Ile Gly Ala Cys Phe Leu Pro Leu Val Trp Ile Val
 20 25 30
 Asn Thr Phe Trp Phe Phe Ser Asp Ala Phe Cys Lys Pro Ile Asn Ala
 10 35 40 45
 His Arg Arg Gln Ile Arg Lys Tyr Val Ile Ala Ser Ile Val Gly Ser
 50 55 60
 Ile Phe Trp Ile Ile Val Leu Ser Ala Trp Glu Ile Phe Phe Gln His
 65 70 75 80
 15 Tyr Arg Ala Gln Gly Leu Val Trp Thr Asp Phe Leu Thr Phe Val Phe
 85 90 95
 Pro Thr Gly Arg Val
 100

20 <210> 8

<211> 101

<212> PRT

<213> *Drosophila melanogaster*

<400> 8

25 Met Asn Ile Ser Lys Ala Pro Asn Pro Arg Lys Leu Glu Leu Cys Arg
 1 5 10 15
 Lys Tyr Phe Phe Ala Gly Phe Ala Phe Leu Pro Phe Val Trp Ala Ile
 20 25 30
 Asn Val Cys Trp Phe Phe Thr Glu Ala Phe His Lys Pro Pro Phe Ser
 30 35 40 45
 Glu His Ser Gln Ile Lys Arg Tyr Val Ile Tyr Ser Ala Val Gly Thr
 50 55 60
 Leu Phe Trp Leu Ile Val Leu Thr Ala Trp Ile Ile Ile Phe Gln Thr
 65 70 75 80
 35 Asn Arg Thr Ala Trp Gly Ala Thr Ala Asp Tyr Met Ser Phe Ile Ile
 85 90 95
 Pro Leu Gly Ser Ala
 100

<210> 9
 <211> 101
 <212> PRT
 <213> rat
 <400> 9

5 Met Asn Leu Glu Arg Val Ser Asn Glu Glu Lys Leu Asn Leu Cys Arg
 1 5 10 15
 Lys Tyr Tyr Leu Gly Gly Phe Ala Phe Leu Pro Phe Leu Trp Leu Val
 20 25 30
 Asn Ile Phe Trp Phe Phe Lys Glu Ala Phe Phe Ala Pro Ala Tyr Ser
 10 35 40 45
 Glu Gln Ser Gln Ile Lys Gly Tyr Val Trp Arg Ser Ala Val Gly Phe
 50 55 60
 Leu Phe Trp Val Ile Val Leu Thr Thr Trp Ile Thr Ile Phe Gln Ile
 65 70 75 80
 15 Tyr Arg Pro Arg Trp Gly Ala Leu Gly Asp Tyr Leu Ser Phe Thr Ile
 85 90 95
 Pro Leu Gly Thr Pro
 100

20 <210> 10
 <211> 101
 <212> PRT
 <213> mouse
 <400> 10

25 Met Asn Leu Glu Arg Val Ser Asn Glu Glu Lys Leu Asn Leu Cys Arg
 1 5 10 15
 Lys Tyr Tyr Leu Gly Gly Phe Ala Phe Leu Pro Phe Leu Trp Leu Val
 20 25 30
 Asn Ile Phe Trp Phe Phe Arg Glu Ala Phe Leu Ala Pro Ala Tyr Thr
 30 35 40 45
 Glu Gln Ser Gln Ile Lys Gly Tyr Val Trp Arg Ser Ala Val Gly Phe
 50 55 60
 Leu Phe Trp Val Ile Ile Leu Ala Thr Trp Ile Thr Ile Phe Gln Ile
 65 70 75 80
 35 Tyr Arg Pro Arg Trp Gly Ala Leu Gly Asp Tyr Leu Ser Phe Thr Ile
 85 90 95
 Pro Leu Gly Thr Pro
 100

<210> 11
 <211> 71
 <212> PRT
 <213> Bos taurus
 <400> 11

5 Met Asn Leu Glu Arg Val Ser Asn Glu Glu Lys Leu Asn Leu Cys Arg
 1 5 10 15
 Lys Tyr Tyr Leu Gly Gly Phe Ala Phe Leu Pro Phe Leu Trp Leu Val
 20 25 30
 Asn Ile Phe Trp Phe Phe Arg Glu Ala Phe Ile Val Pro Ala Tyr Thr
 10 35 40 45
 Glu Gln Ser Gln Ile Lys Gly Tyr Val Trp Arg Ser Ala Val Gly Phe
 50 55 60
 Phe Leu Trp Val Ile Val Leu
 65 70

15
 <210> 12
 <211> 101
 <212> PRT
 <213> human
 20 <400> 12

Met Asn Leu Glu Arg Val Ser Asn Glu Glu Lys Leu Asn Leu Cys Arg
 1 5 10 15
 Lys Tyr Tyr Leu Gly Gly Phe Ala Phe Leu Pro Phe Leu Trp Leu Val
 20 25 30
 25 Asn Ile Phe Trp Phe Phe Arg Glu Ala Phe Leu Val Pro Ala Tyr Thr
 35 40 45
 Glu Gln Ser Gln Ile Lys Gly Tyr Val Trp Arg Ser Ala Val Gly Phe
 50 55 60
 Leu Phe Trp Val Ile Val Leu Thr Ser Trp Ile Thr Ile Phe Gln Ile
 30 65 70 75 80
 Tyr Arg Pro Arg Trp Gly Ala Leu Gly Asp Tyr Leu Ser Phe Thr Ile
 85 90 95
 Pro Leu Gly Thr Pro
 100

35
 <210> 13
 <211> 721
 <212> PRT

<213> *Caenorhabditis elegans*
 <400> 13
 Met Lys Lys Trp Leu Val Ile Val Leu Ile Ile Ala Gly Ile Arg Cys
 1 5 10 15
 Asp Gly Phe Ser Asp Gln Val Phe Arg Thr Leu Phe Ile Gly Glu Gly
 5 20 25 30
 Asn Ala Cys Tyr Arg Thr Phe Asn Lys Thr His Glu Phe Gly Cys Gln
 35 40 45
 Ala Asn Arg Glu Asn Glu Asn Gly Leu Ile Val Arg Ile Asp Lys Gln
 50 55 60
 10 Glu Asp Phe Lys Asn Leu Asp Ser Cys Trp Asn Ser Phe Tyr Pro Lys
 65 70 75 80
 Tyr Ser Gly Lys Tyr Trp Ala Leu Leu Pro Val Asn Leu Ile Arg Arg
 85 90 95
 Asp Thr Ile Ser Gln Leu Lys Ser Ser Lys Cys Leu Ser Gly Ile Val
 15 100 105 110
 Leu Tyr Asn Ser Gly Glu Ser Ile His Pro Gly Asp Glu Ser Thr Ala
 115 120 125
 Ala Ser His Asp Ala Glu Cys Pro Asn Ala Ala Ser Asp Tyr Tyr Leu
 130 135 140
 20 Gln Asp Lys Asn Glu Glu Tyr Cys Glu Arg Lys Ile Asn Ser Arg Gly
 145 150 155 160
 Ala Ile Thr Arg Asp Gly Leu Met Lys Ile Asp Trp Arg Ile Gln Met
 165 170 175
 Val Phe Ile Asp Asn Ser Thr Asp Leu Glu Ile Ile Glu Lys Cys Tyr
 25 180 185 190
 Ser Met Phe Asn Lys Pro Lys Glu Asp Gly Ser Ser Gly Tyr Pro Tyr
 195 200 205
 Cys Gly Met Ser Phe Arg Leu Ala Asn Met Ala Ala Gly Asn Ser Glu
 210 215 220
 30 Ile Cys Tyr Arg Arg Gly Lys Asn Asp Ala Lys Leu Phe Gln Met Asn
 225 230 235 240
 Ile Asp Ser Gly Asp Ala Pro Gln Leu Cys Gly Ala Met His Ser Asp
 245 250 255
 Asn Ile Phe Ala Phe Pro Thr Pro Ile Pro Thr Ser Pro Thr Asn Glu
 35 260 265 270
 Thr Ile Ile Thr Ser Lys Tyr Met Met Val Thr Ala Arg Met Asp Ser
 275 280 285
 Phe Gly Met Ile Pro Glu Ile Ser Val Gly Glu Val Ser Val Leu Thr

	290	295	300
	Ser Ile Ile Ser Val Leu Ala Ala Ala Arg Ser Met Gly Thr Gln Ile		
	305	310	315 320
	Glu Lys Trp Gln Lys Ala Ser Asn Thr Ser Asn Arg Asn Val Phe Phe		
	325	330	335
5	Ala Phe Phe Asn Gly Glu Ser Leu Asp Tyr Ile Gly Ser Gly Ala Ala		
	340	345	350
	Ala Tyr Gln Met Glu Asn Gly Lys Phe Pro Gln Met Ile Arg Ser Asp		
	355	360	365
	Arg Thr His Ile His Pro Ile Arg Pro Asn Glu Leu Asp Tyr Ile Leu		
10	370	375	380
	Glu Val Gln Gln Ile Gly Val Ala Lys Gly Arg Lys Tyr Tyr Val His		
	385	390	395 400
	Val Asp Gly Glu Arg Tyr Gln Gln Asn Lys Thr Gln Thr Asp Arg Val		
	405	410	415
15	Ile Asp Arg Ile Glu Arg Gly Leu Arg Ser His Ala Phe Asp Leu Glu		
	420	425	430
	Lys Pro Ser Gly Ser Gly Asp Arg Val Pro Pro Ala Ser Trp His Ser		
	435	440	445
	Phe Ala Lys Ala Asp Ala His Val Gln Ser Val Leu Leu Ala Pro Tyr		
20	450	455	460
	Gly Lys Glu Tyr Glu Tyr Gln Arg Val Asn Ser Ile Leu Asp Lys Asn		
	465	470	475 480
	Glu Trp Thr Glu Asp Glu Arg Glu Lys Ala Ile Gln Glu Ile Glu Ala		
	485	490	495
25	Val Ser Thr Ala Ile Leu Ala Ala Ala Ala Asp Tyr Val Gly Val Glu		
	500	505	510
	Thr Asp Glu Val Val Ala Lys Val Asp Lys Lys Leu Ile Thr Thr Ile		
	515	520	525
	Phe Asp Cys Leu Ile Thr Ser Asn Phe Trp Phe Asp Cys Asp Phe Met		
30	530	535	540
	Gln Lys Leu Asp Gly Gly Arg Tyr His Lys Leu Phe Asn Ser Tyr Gly		
	545	550	555 560
	Phe Asn Gln Lys Ser Thr Tyr Ile Ser Met Glu Ser His Thr Ala Phe		
	565	570	575
35	Pro Thr Val Leu His Trp Leu Thr Ile Phe Ala Leu Gly Ser Asp Lys		
	580	585	590
	Glu Thr Leu Asn Val Lys Ser Glu Lys Ser Cys Ser His Leu Gly Gln		
	595	600	605

Phe Gln Ala Met Tyr Thr Tyr Thr Trp Gln Pro Asn Pro Tyr Thr Gly
 610 615 620
 Asn Phe Ser Cys Leu Lys Ser Ala Ile Val Lys Lys Val Met Val Ser
 625 630 635 640
 Pro Ala Val Asp Ser Gln Thr Pro Glu Glu Glu Met Asn Thr Arg Tyr
 5 645 650 655
 Ser Thr Trp Met Glu Ser Val Tyr Ile Ile Glu Ser Val Asn Leu Tyr
 660 665 670
 Leu Met Glu Asp Ala Ser Phe Glu Tyr Thr Met Ile Leu Ile Ala Val
 675 680 685
 10 Ile Ser Ala Leu Leu Ser Ile Phe Ala Val Gly Arg Cys Ser Glu Thr
 690 695 700
 Thr Phe Ile Val Asp Glu Gly Glu Pro Ala Ala Glu Gly Gly Glu Pro
 705 710 715 720
 Leu
 15
 <210> 14
 <211> 708
 <212> PRT
 <213> *Drosophila melanogaster*
 20 <400> 14
 Ala Thr Ala Gly Gly Gly Ser Gly Ala Asp Pro Gly Ser Arg Gly Leu
 1 5 10 15
 Leu Arg Leu Leu Ser Phe Cys Val Leu Leu Ala Gly Leu Cys Arg Gly
 20 25 30
 25 Asn Ser Val Glu Arg Lys Ile Tyr Ile Pro Leu Asn Lys Thr Ala Pro
 35 40 45
 Cys Val Arg Leu Leu Asn Ala Thr His Gln Ile Gly Cys Gln Ser Ser
 50 55 60
 Ile Ser Gly Asp Thr Gly Val Ile His Val Val Glu Lys Glu Glu Asp
 65 70 75 80
 30 Leu Gln Trp Val Leu Thr Asp Gly Pro Asn Pro Pro Tyr Met Val Leu
 85 90 95
 Leu Glu Ser Lys His Phe Thr Arg Asp Leu Met Glu Lys Leu Lys Gly
 100 105 110
 35 Arg Thr Ser Arg Ile Ala Gly Leu Ala Val Ser Leu Thr Lys Pro Ser
 115 120 125
 Pro Ala Ser Gly Phe Ser Pro Ser Val Gln Cys Pro Asn Asp Gly Phe
 130 135 140

Gly Val Tyr Ser Asn Ser Tyr Gly Pro Glu Phe Ala His Cys Arg Glu
 145 150 155 160
 Ile Gln Trp Asn Ser Leu Gly Asn Gly Leu Ala Tyr Glu Asp Phe Ser
 165 170 175
 Phe Pro Ile Phe Leu Leu Glu Asp Glu Asn Glu Thr Lys Val Ile Lys
 5 180 185 190
 Gln Cys Tyr Gln Asp His Asn Leu Ser Gln Asn Gly Ser Ala Pro Thr
 195 200 205
 Phe Pro Leu Cys Ala Met Gln Leu Phe Ser His Met His Ala Val Ile
 210 215 220
 10 Ser Thr Ala Thr Cys Met Arg Arg Ser Ser Ile Gln Ser Thr Phe Ser
 225 230 235 240
 Ile Asn Pro Glu Ile Val Cys Asp Pro Leu Ser Asp Tyr Asn Val Trp
 245 250 255
 Ser Met Leu Lys Pro Ile Asn Thr Thr Gly Thr Leu Lys Pro Asp Asp
 15 260 265 270
 Arg Val Val Val Ala Ala Thr Arg Leu Asp Ser Arg Ser Phe Phe Trp
 275 280 285
 Asn Val Ala Pro Gly Ala Glu Ser Ala Val Ala Ser Phe Val Thr Gln
 290 295 300
 20 Leu Ala Ala Ala Glu Ala Leu Gln Lys Ala Pro Asp Val Thr Thr Leu
 305 310 315 320
 Pro Arg Asn Val Met Phe Val Phe Phe Gln Gly Glu Thr Phe Asp Tyr
 325 330 335
 25 Ile Gly Ser Ser Arg Met Val Tyr Asp Met Glu Lys Gly Lys Phe Pro
 340 345 350
 Val Gln Leu Glu Asn Val Asp Ser Phe Val Glu Leu Gly Gln Val Ala
 355 360 365
 Leu Arg Thr Ser Leu Glu Leu Trp Met His Thr Asp Pro Val Ser Gln
 30 370 375 380
 Lys Asn Glu Ser Val Arg Asn Gln Val Glu Asp Leu Leu Ala Thr Leu
 385 390 395 400
 Glu Lys Ser Gly Ala Gly Val Pro Ala Val Ile Leu Arg Arg Pro Asn
 405 410 415
 35 Gln Ser Gln Pro Leu Pro Pro Ser Ser Leu Gln Arg Phe Leu Arg Ala
 420 425 430
 Arg Asn Ile Ser Gly Val Val Leu Ala Asp His Ser Gly Ala Phe His
 435 440 445

Asn Lys Tyr Tyr Gln Ser Ile Tyr Asp Thr Ala Glu Asn Ile Asn Val
 450 455 460
 Ser Tyr Pro Glu Trp Leu Ser Pro Glu Glu Asp Leu Asn Phe Val Thr
 465 470 475 480
 Asp Thr Ala Lys Ala Leu Ala Asp Val Ala Thr Val Leu Gly Arg Ala
 5 485 490 495
 Leu Tyr Glu Leu Ala Gly Gly Thr Asn Phe Ser Asp Thr Val Gln Ala
 500 505 510
 Asp Pro Gln Thr Val Thr Arg Leu Leu Tyr Gly Phe Leu Ile Lys Ala
 515 520 525
 10 Asn Asn Ser Trp Phe Gln Ser Ile Leu Arg Gln Asp Leu Arg Ser Tyr
 530 535 540
 Leu Gly Asp Gly Pro Leu Gln His Tyr Ile Ala Val Ser Ser Pro Thr
 545 550 555 560
 Asn Thr Thr Tyr Val Val Gln Tyr Ala Leu Ala Asn Leu Thr Gly Thr
 15 565 570 575
 Val Val Asn Leu Thr Arg Glu Gln Cys Gln Asp Pro Ser Lys Val Pro
 580 585 590
 Ser Glu Asn Lys Asp Leu Tyr Glu Tyr Ser Trp Val Gln Gly Pro Leu
 595 600 605
 20 His Ser Asn Glu Thr Asp Arg Leu Pro Arg Cys Val Arg Ser Thr Ala
 610 615 620
 Arg Leu Ala Arg Ala Leu Ser Pro Ala Phe Glu Leu Ser Gln Trp Ser
 625 630 635 640
 Ser Thr Glu Tyr Ser Thr Trp Thr Glu Ser Arg Trp Lys Asp Ile Arg
 25 645 650 655
 Ala Arg Ile Phe Leu Ile Ala Ser Lys Glu Leu Glu Leu Ile Thr Leu
 660 665 670
 Thr Val Gly Phe Gly Ile Leu Ile Phe Ser Leu Ile Val Thr Tyr Cys
 675 680 685
 30 Ile Asn Ala Lys Ala Asp Val Leu Phe Ile Ala Pro Arg Glu Pro Gly
 690 695 700
 Ala Val Ser Tyr
 705
 35 <210> 15
 <211> 716
 <212> PRT
 <213> human

<400> 15

His Glu Pro Lys Arg Ser His Ala Thr Leu Gln Phe Leu Asp Ala Ile
 1 5 10 15
 Ser Trp Glu Ser Ser Met Glu Met Arg Leu Asn Ala Ala Ser Ile Trp
 20 25 30
 5 Leu Leu Ile Leu Ser Tyr Gly Ala Thr Ile Ala Gln Gly Glu Arg Thr
 35 40 45
 Arg Asp Lys Met Tyr Glu Pro Ile Gly Gly Ala Ser Cys Phe Arg Arg
 50 55 60
 Leu Asn Gly Thr His Gln Thr Gly Cys Ser Ser Thr Tyr Ser Gly Ser
 10 65 70 75 80
 Val Gly Val Leu His Leu Ile Asn Val Glu Ala Asp Leu Glu Phe Leu
 85 90 95
 Leu Ser Ser Pro Pro Ser Pro Pro Tyr Ala Pro Met Ile Pro Pro His
 100 105 110
 15 Leu Phe Thr Arg Asn Asn Leu Met Arg Leu Lys Glu Ala Gly Pro Lys
 115 120 125
 Asn Ile Ser Val Val Leu Leu Ile Asn Arg Thr Asn Gln Met Lys Gln
 130 135 140
 Phe Ser His Glu Leu Asn Cys Pro Asn Gln Tyr Ser Gly Leu Asn Ser
 145 150 155 160
 20 Thr Ser Glu Thr Cys Asp Ala Ser Asn Pro Ala Lys Asn Trp Asn Pro
 165 170 175
 Trp Gly Thr Gly Leu Leu His Glu Asp Phe Pro Phe Pro Ile Tyr Tyr
 180 185 190
 25 Ile Ala Asp Leu Asp Gln Val Thr Lys Leu Glu Lys Cys Phe Gln Asp
 195 200 205
 Phe Asn Asn His Asn Tyr Glu Thr His Ala Leu Arg Ser Leu Cys Ala
 210 215 220
 Val Glu Val Lys Ser Phe Met Ser Ala Ala Val Asn Thr Glu Val Cys
 225 230 235 240
 30 Met Arg Arg Thr Asn Phe Ile Asn Asn Leu Gly Gly Ser Lys Tyr Cys
 245 250 255
 Asp Pro Leu Glu Gly Arg Asn Val Tyr Ala Thr Leu Tyr Pro Glu Ser
 260 265 270
 35 Gln Gln Ser Lys Thr Thr Trp Arg Gln Ser Ile Arg Met Lys Ser Ser
 275 280 285
 Ile Ser Asn Leu Ser Pro Gly His His His His Val Arg Trp Arg Arg
 290 295 300

Ser Trp Ser His Gly Leu Pro Tyr Gly Ile Cys Trp Phe Gln Leu Ser
 305 310 315 320
 Val Gly Tyr Leu Leu Lys Gln Leu Leu Pro Pro Gln Ser Lys Asp Leu
 325 330 335
 His Asn Val Leu Phe Val Thr Phe Asn Gly Glu Ser Tyr Asp Tyr Ile
 5 340 345 350
 Gly Ser Gln Arg Phe Val Tyr Asp Met Glu Lys Leu Gln Phe Pro Thr
 355 360 365
 Glu Ser Thr Gly Thr Pro Pro Ile Ala Phe Asp Asn Ile Asp Phe Met
 370 375 380
 10 Leu Asp Ile Gly Thr Leu Asp Asp Ile Ser Asn Ile Lys Leu His Ala
 385 390 395 400
 Leu Asn Gly Thr Thr Leu Ala Gln Gln Ile Leu Glu Arg Leu Asn Asn
 405 410 415
 15 Tyr Ala Lys Ser Pro Arg Tyr Gly Phe Asn Leu Asn Ile Gln Ser Glu
 420 425 430
 Met Ser Ala His Leu Pro Pro Thr Ser Ala Gln Ser Phe Leu Arg Arg
 435 440 445
 Asp Pro Asn Phe Asn Ala Leu Ile Leu Asn Ala Arg Pro Thr Asn Lys
 20 450 455 460
 Tyr Tyr His Ser Ile Tyr Asp Asp Ala Asp Asn Val Asp Phe Thr Tyr
 465 470 475 480
 Ala Asn Thr Ser Lys Asp Phe Thr Gln Leu Thr Glu Val Asn Asp Phe
 485 490 495
 25 Lys Ser Leu Asn Pro Asp Ser Leu Gln Met Lys Val Arg Asn Val Ser
 500 505 510
 Ser Ile Val Ala Met Ala Leu Tyr Gln Thr Ile Thr Gly Lys Glu Tyr
 515 520 525
 Thr Gly Thr Lys Val Ala Asn Pro Leu Met Ala Asp Glu Phe Leu Tyr
 30 530 535 540
 Cys Phe Leu Gln Ser Ala Asp Cys Pro Leu Phe Lys Ala Ala Ser Tyr
 545 550 555 560
 Pro Gly Ser Gln Leu Thr Asn Leu Pro Pro Met Arg Tyr Ile Ser Val
 565 570 575
 35 Leu Gly Gly Ser Gln Glu Ser Ser Gly Tyr Thr Tyr Arg Leu Leu Gly
 580 585 590
 Tyr Leu Leu Ser Gln Leu Gln Pro Asp Ile His Arg Asp Asn Cys Thr
 595 600 605

Asp Leu Pro Leu His Tyr Phe Ala Gly Phe Asn Asn Ile Gly Glu Cys
 610 615 620
 Arg Leu Thr Thr Gln Asn Tyr Ser His Ala Leu Ser Pro Ala Phe Leu
 625 630 635 640
 Ile Asp Gly Tyr Asp Trp Ser Ser Gly Met Tyr Ser Thr Trp Ala Glu
 5 645 650 655
 Ser Thr Trp Ser Gln Phe Ser Ala Arg Ile Phe Leu Arg Pro Ser Asn
 660 665 670
 Val His Gln Val Thr Thr Leu Ser Val Gly Ile Val Val Leu Ile Ile
 675 680 685
 10 Ser Phe Cys Leu Val Tyr Ile Ile Ser Ser Arg Ser Glu Val Leu Phe
 690 695 700
 Glu Asp Leu Pro Ala Ser Asn Ala Ala Leu Phe Gly
 705 710 715

 15 <210> 16
 <211> 257
 <212> PRT
 <213> Artificial Sequence
 <220>
 20 <223> Description of Artificial Sequence: Synthetic
 Sequence
 <400> 16
 Met Ala Ala Ala Val Phe Phe Gly Cys Ala Phe Asp Ala Phe Gly Pro
 1 5 10 15
 25 Ala Leu Ala Leu Tyr Glu Phe Thr Ile Ala Thr Glu Pro Leu Arg Phe
 20 25 30
 Ile Phe Leu Ile Ala Gly Ala Phe Phe Gly Leu Val Ser Leu Leu Ile
 35 40 45
 Ser Ser Leu His Trp Phe Met Ala Arg Val Ile Ile Asp Ile Lys Asp
 30 50 55 60
 Gly Pro Thr Gln Lys Tyr Leu Lys Ile Phe Gly Ala Phe Val Ser Val
 65 70 75 80
 Tyr Leu Gln Glu Met Phe Arg Phe Ala Tyr Tyr Met Leu Leu Lys Lys
 85 90 95
 35 Ala Ser Glu Gly Leu Asn Ser Ile Asn Pro Gly Glu Thr Ala Pro Gln
 100 105 110
 Met Arg Leu Leu Ala Tyr Val Ser Gly Arg Gly Phe Gly Ile Met Ser
 115 120 125

Gly Val Phe Thr Phe Val Asn Thr Leu Ser Asp Ser Leu Val Pro Gly
 130 135 140
 Thr Val Gly Ile His Gly Asp Trp Pro Gln Phe Phe Leu Tyr Ser Ala
 145 150 155 160
 Phe Tyr Thr Leu Val Ile Ile Leu Leu His Val Ala Trp Gly Ile Val
 5 165 170 175
 Phe Phe Asp Gly Cys Asp Lys Lys Lys Trp Gly Ile Leu Leu Ile Glu
 180 185 190
 Leu Leu Thr His Leu Leu Val Ser Ala Phe Thr Phe Ile Ser Ser Tyr
 195 200 205
 10 Tyr Gly Ile Gly Leu Ala Ser Ala Phe Ile Ile Leu Val His Met Gly
 210 215 220
 Thr Trp Ala Phe Leu Ala Ala Ile Gly Ser Cys Arg Ser Leu Lys Leu
 225 230 235 240
 Cys Lys Leu Cys Gln Asp Lys Asn Phe Leu Leu Leu Asn Gln Arg Ser
 15 245 250 255
 Arg

 <210> 17
 <211> 257
 20 <212> PRT
 <213> Artificial Sequence
 <220>
 <223> Description of Artificial Sequence: Synthetic
 Sequence
 25 <400> 17
 Met Thr Asp Ala Val Phe Phe Gly Cys Ala Phe Ile Glu Phe Gly Pro
 1 5 10 15
 Ala Leu Ala Leu Tyr Val Gly Thr Ile Ala Thr Glu Pro Leu Arg Ile
 20 25 30
 30 His Phe Leu Ile Ala Gly Ala Phe Phe Trp Ile Val Ser Leu Leu Ile
 35 40 45
 Ser Ser Leu Val Lys Phe Met Ala Arg Val Ile Ile Asp Asn Leu Asp
 50 55 60
 Gly Pro Thr Gln Lys Tyr Leu Leu Met Phe Gly Ala Phe Val Ser Val
 35 65 70 75 80
 Tyr Ile Asn Glu Met Phe Arg Phe Ala Tyr Tyr Lys Gln Leu Lys Lys
 85 90 95
 Ala Ser Glu Gly Leu Lys Arg Ile Asn Pro Gly Glu Thr Ala Pro Ser

100 105 110
 Ser Arg Leu Leu Ala Tyr Val Ser Gly Leu Thr Phe Gly Ile Met Ser
 115 120 125
 Gly Val Phe Ser Val Val Asn Thr Leu Ser Asp Ser Leu Gly Trp Gly
 130 135 140
 5 Thr Val Gly Ile His Gly Asp Ser Tyr Gln Phe Phe Leu Tyr Ser Ala
 145 150 155 160
 Phe Met Ala Leu Val Ile Ile Leu Leu His Val Phe Asp Gly Ile Val
 165 170 175
 Phe Phe Asp Gly Cys Glu Glu Lys Lys Trp Gly Ile Leu Leu Ile Val
 10 180 185 190
 Phe Leu Thr His Leu Leu Val Ser Ala Gln Gly Phe Ile Ser Ser Tyr
 195 200 205
 Tyr Gly Ile Asn His Ala Ser Ala Phe Ile Ile Leu Val Leu Ile Gly
 210 215 220
 15 Thr Trp Ala Phe Leu Ala Ala Gly Lys Ser Cys Arg Ser Leu Lys Leu
 225 230 235 240
 Cys Leu Met Cys Gln Asp Lys Asn Phe Leu Leu Tyr Gln Gln Arg Ser
 245 250 255
 Arg
 20
 <210> 18
 <211> 257
 <212> PRT
 <213> Artificial Sequence
 25 <220>
 <223> Description of Artificial Sequence: Synthetic
 Sequence
 <400> 18
 Met Thr Ala Asp Val Phe Phe Gly Cys Ala Phe Ile Ala Glu Gly Pro
 30 1 5 10 15
 Ala Leu Ala Leu Tyr Val Phe Phe Ile Ala Thr Glu Pro Leu Arg Ile
 20 25 30
 Ile Gly Leu Ile Ala Gly Ala Phe Phe Trp Leu His Ser Leu Leu Ile
 35 35 40 45
 Ser Ser Leu Val Trp Ile Met Ala Arg Val Ile Ile Asp Asn Lys Lys
 50 55 60
 Gly Pro Thr Gln Lys Tyr Leu Leu Ile Leu Gly Ala Phe Val Ser Val
 65 70 75 80

Tyr Ile Gln Met Met Phe Arg Phe Ala Tyr Tyr Lys Leu Asn Lys Lys
 85 90 95
 Ala Ser Glu Gly Leu Lys Ser Gln Asn Pro Gly Glu Thr Ala Pro Ser
 100 105 110
 Met Ser Leu Leu Ala Tyr Val Ser Gly Leu Gly Thr Gly Ile Met Ser
 5 115 120 125
 Gly Val Phe Ser Phe Trp Asn Thr Leu Ser Asp Ser Leu Gly Pro Tyr
 130 135 140
 Thr Val Gly Ile His Gly Asp Ser Pro Ala Phe Phe Leu Tyr Ser Ala
 145 150 155 160
 10 Phe Met Thr Asp Val Ile Ile Leu Leu His Val Phe Trp Glu Ile Val
 165 170 175
 Phe Phe Asp Gly Cys Glu Lys Phe Lys Trp Gly Ile Leu Leu Ile Val
 180 185 190
 Leu Gly Thr His Leu Leu Val Ser Ala Gln Thr His Ile Ser Ser Tyr
 15 195 200 205
 Tyr Gly Ile Asn Leu Ile Ser Ala Phe Ile Ile Leu Val Leu Met Lys
 210 215 220
 Thr Trp Ala Phe Leu Ala Ala Gly Gly Leu Cys Arg Ser Leu Lys Leu
 225 230 235 240
 20 Cys Leu Leu Met Gln Asp Lys Asn Phe Leu Leu Tyr Asn Asn Arg Ser
 245 250 255
 Arg

25 <210> 19
 <211> 257
 <212> PRT
 <213> Artificial Sequence
 <220>

30 <223> Description of Artificial Sequence: Synthetic
 Sequence

<400> 19
 Met Thr Ala Ala Ala Phe Phe Gly Cys Ala Phe Ile Ala Phe Asp Pro
 1 5 10 15
 Ala Leu Ala Leu Tyr Val Phe Thr Glu Ala Thr Glu Pro Leu Arg Ile
 35 20 25 30
 Ile Phe Phe Ile Ala Gly Ala Phe Phe Trp Leu Val Gly Leu Leu Ile
 35 40 45
 Ser Ser Leu Val Trp Phe His Ala Arg Val Ile Ile Asp Asn Lys Asp

50 55 60
 Ile Pro Thr Gln Lys Tyr Leu Leu Ile Phe Lys Ala Phe Val Ser Val
 65 70 75 80
 Tyr Ile Gln Glu Leu Phe Arg Phe Ala Tyr Tyr Lys Leu Leu Met Lys
 85 90 95
 5 Ala Ser Glu Gly Leu Lys Ser Ile Gln Pro Gly Glu Thr Ala Pro Ser
 100 105 110
 Met Arg Arg Leu Ala Tyr Val Ser Gly Leu Gly Phe Ser Ile Met Ser
 115 120 125
 Gly Val Phe Ser Phe Val Thr Thr Leu Ser Asp Ser Leu Gly Pro Gly
 10 130 135 140
 Val Val Gly Ile His Gly Asp Ser Pro Gln Trp Phe Leu Tyr Ser Ala
 145 150 155 160
 Phe Met Thr Leu Tyr Ile Ile Leu Leu His Val Phe Trp Gly Ala Val
 165 170 175
 15 Phe Phe Asp Gly Cys Glu Lys Lys Asp Trp Gly Ile Leu Leu Ile Val
 180 185 190
 Leu Leu Glu His Leu Leu Val Ser Ala Gln Thr Phe Phe Ser Ser Tyr
 195 200 205
 Tyr Gly Ile Asn Leu Ala Gly Ala Phe Ile Ile Leu Val Leu Met Gly
 20 210 215 220
 His Trp Ala Phe Leu Ala Ala Gly Gly Ser Ile Arg Ser Leu Lys Leu
 225 230 235 240
 Cys Leu Leu Cys Lys Asp Lys Asn Phe Leu Leu Tyr Asn Gln Leu Ser
 245 250 255
 25 Arg

 <210> 20
 <211> 257
 <212> PRT
 30 <213> Artificial Sequence
 <220>
 <223> Description of Artificial Sequence: Synthetic
 Sequence
 <400> 20
 35 Met Thr Ala Ala Val Ala Phe Gly Cys Ala Phe Ile Ala Phe Gly Asp
 1 5 10 15
 Ala Leu Ala Leu Tyr Val Phe Thr Ile Glu Thr Glu Pro Leu Arg Ile
 20 25 30

Ile Phe Leu Phe Ala Gly Ala Phe Phe Trp Leu Val Ser Gly Leu Ile
 35 40 45
 Ser Ser Leu Val Trp Phe Met His Arg Val Ile Ile Asp Asn Lys Asp
 50 55 60
 Gly Ile Thr Gln Lys Tyr Leu Leu Ile Phe Gly Lys Phe Val Ser Val
 5 65 70 75 80
 Tyr Ile Gln Glu Met Leu Arg Phe Ala Tyr Tyr Lys Leu Leu Lys Met
 85 90 95
 Ala Ser Glu Gly Leu Lys Ser Ile Asn Asn Gly Glu Thr Ala Pro Ser
 100 105 110
 10 Met Arg Leu Gln Ala Tyr Val Ser Gly Leu Gly Phe Gly Arg Met Ser
 115 120 125
 Gly Val Phe Ser Phe Val Asn Ser Leu Ser Asp Ser Leu Gly Pro Gly
 130 135 140
 Thr Thr Gly Ile His Gly Asp Ser Pro Gln Phe Val Leu Tyr Ser Ala
 15 145 150 155 160
 Phe Met Thr Leu Val Trp Ile Leu Leu His Val Phe Trp Gly Ile Tyr
 165 170 175
 Phe Phe Asp Gly Cys Glu Lys Lys Lys Ala Gly Ile Leu Leu Ile Val
 180 185 190
 20 Leu Leu Thr Asp Leu Leu Val Ser Ala Gln Thr Phe Ile Glu Ser Tyr
 195 200 205
 Tyr Gly Ile Asn Leu Ala Ser Phe Phe Ile Ile Leu Val Leu Met Gly
 210 215 220
 Thr Gly Ala Phe Leu Ala Ala Gly Gly Ser Cys His Ser Leu Lys Leu
 25 225 230 235 240
 Cys Leu Leu Cys Gln Ile Lys Asn Phe Leu Leu Tyr Asn Gln Arg Lys
 245 250 255
 Arg
 30 <210> 21
 <211> 257
 <212> PRT
 <213> Artificial Sequence
 <220>
 35 <223> Description of Artificial Sequence: Synthetic
 Sequence
 <400> 21
 Met Thr Ala Ala Val Phe Ala Gly Cys Ala Phe Ile Ala Phe Gly Pro

1 5 10 15
 Asp Leu Ala Leu Tyr Val Phe Thr Ile Ala Glu Glu Pro Leu Arg Ile
 20 25 30
 Ile Phe Leu Ile Phe Gly Ala Phe Phe Trp Leu Val Ser Leu Gly Ile
 35 40 45
 5 Ser Ser Leu Val Trp Phe Met Ala His Val Ile Ile Asp Asn Lys Asp
 50 55 60
 Gly Pro Ile Gln Lys Tyr Leu Leu Ile Phe Gly Ala Lys Val Ser Val
 65 70 75 80
 Tyr Ile Gln Glu Met Phe Leu Phe Ala Tyr Tyr Lys Leu Leu Lys Lys
 10 85 90 95
 Met Ser Glu Gly Leu Lys Ser Ile Asn Pro Asn Glu Thr Ala Pro Ser
 100 105 110
 Met Arg Leu Leu Gln Tyr Val Ser Gly Leu Gly Phe Gly Ile Arg Ser
 115 120 125
 15 Gly Val Phe Ser Phe Val Asn Thr Ser Ser Asp Ser Leu Gly Pro Gly
 130 135 140
 Thr Val Thr Ile His Gly Asp Ser Pro Gln Phe Phe Val Tyr Ser Ala
 145 150 155 160
 Phe Met Thr Leu Val Ile Trp Leu Leu His Val Phe Trp Gly Ile Val
 20 165 170 175
 Tyr Phe Asp Gly Cys Glu Lys Lys Lys Trp Ala Ile Leu Leu Ile Val
 180 185 190
 Leu Leu Thr His Asp Leu Val Ser Ala Gln Thr Phe Ile Ser Glu Tyr
 195 200 205
 25 Tyr Gly Ile Asn Leu Ala Ser Ala Gly Ile Ile Leu Val Leu Met Gly
 210 215 220
 Thr Trp His Phe Leu Ala Ala Gly Gly Ser Cys Arg Ile Leu Lys Leu
 225 230 235 240
 Cys Leu Leu Cys Gln Asp Leu Asn Phe Leu Leu Tyr Asn Gln Arg Ser
 30 245 250 255
 Met

35 <210> 22
 <211> 257
 <212> PRT
 <213> Artificial Sequence
 <220>
 <223> Description of Artificial Sequence: Synthetic

Sequence

<400> 22

Met Thr Ala Ala Val Phe Phe Ala Cys Ala Phe Ile Ala Phe Gly Pro
 1 5 10 15
 Ala Asp Ala Leu Tyr Val Phe Thr Ile Ala Thr Phe Pro Leu Arg Ile
 5 20 25 30
 Ile Phe Leu Ile Ala His Ala Phe Phe Trp Leu Val Ser Leu Leu Lys
 35 40 45
 Ser Ser Leu Val Trp Phe Met Ala Arg Leu Ile Ile Asp Asn Lys Asp
 50 55 60
 10 Gly Pro Thr Met Lys Tyr Leu Leu Ile Phe Gly Ala Phe Asn Ser Val
 65 70 75 80
 Tyr Ile Gln Glu Met Phe Arg Gln Ala Tyr Tyr Lys Leu Leu Lys Lys
 85 90 95
 Ala Arg Glu Gly Leu Lys Ser Ile Asn Pro Gly Ser Thr Ala Pro Ser
 15 100 105 110
 Met Arg Leu Leu Ala Thr Val Ser Gly Leu Gly Phe Gly Ile Met Val
 115 120 125
 Gly Val Phe Ser Phe Val Asn Thr Leu Trp Asp Ser Leu Gly Pro Gly
 130 135 140
 20 Thr Val Gly Tyr His Gly Asp Ser Pro Gln Phe Phe Leu Ala Ser Ala
 145 150 155 160
 Phe Met Thr Leu Val Ile Ile Asp Leu His Val Phe Trp Gly Ile Val
 165 170 175
 Phe Glu Asp Gly Cys Glu Lys Lys Lys Trp Gly Phe Leu Leu Ile Val
 25 180 185 190
 Leu Leu Thr His Leu Gly Val Ser Ala Gln Thr Phe Ile Ser Ser His
 195 200 205
 Tyr Gly Ile Asn Leu Ala Ser Ala Phe Lys Ile Leu Val Leu Met Gly
 210 215 220
 30 Thr Trp Ala Leu Leu Ala Ala Gly Gly Ser Cys Arg Ser Met Lys Leu
 225 230 235 240
 Cys Leu Leu Cys Gln Asp Lys Gln Phe Leu Leu Tyr Asn Gln Arg Ser
 245 250 255
 Arg
 35
 <210> 23
 <211> 257
 <212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
Sequence

<400> 23

5 Met Thr Ala Ala Val Phe Phe Gly Ala Ala Phe Ile Ala Phe Gly Pro
 1 5 10 15
 Ala Leu Asp Leu Tyr Val Phe Thr Ile Ala Thr Glu Glu Leu Arg Ile
 20 25 30
 Ile Phe Leu Ile Ala Gly Phe Phe Phe Trp Leu Val Ser Leu Leu Ile
 10 35 40 45
 Gly Ser Leu Val Trp Phe Met Ala Arg Val His Ile Asp Asn Lys Asp
 50 55 60
 Gly Pro Thr Gln Ile Tyr Leu Leu Ile Phe Gly Ala Phe Val Lys Val
 65 70 75 80
 15 Tyr Ile Gln Glu Met Phe Arg Phe Leu Tyr Tyr Lys Leu Leu Lys Lys
 85 90 95
 Ala Ser Met Gly Leu Lys Ser Ile Asn Pro Gly Glu Asn Ala Pro Ser
 100 105 110
 Met Arg Leu Leu Ala Tyr Gln Ser Gly Leu Gly Phe Gly Ile Met Ser
 20 115 120 125
 Arg Val Phe Ser Phe Val Asn Thr Leu Ser Ser Ser Leu Gly Pro Gly
 130 135 140
 Thr Val Gly Ile Thr Gly Asp Ser Pro Gln Phe Phe Leu Tyr Val Ala
 145 150 155 160
 25 Phe Met Thr Leu Val Ile Ile Leu Trp His Val Phe Trp Gly Ile Val
 165 170 175
 Phe Phe Tyr Gly Cys Glu Lys Lys Lys Trp Gly Ile Ala Leu Ile Val
 180 185 190
 Leu Leu Thr His Leu Leu Asp Ser Ala Gln Thr Phe Ile Ser Ser Tyr
 30 195 200 205
 Glu Gly Ile Asn Leu Ala Ser Ala Phe Ile Phe Leu Val Leu Met Gly
 210 215 220
 Thr Trp Ala Phe Gly Ala Ala Gly Gly Ser Cys Arg Ser Leu His Leu
 225 230 235 240
 35 Cys Leu Leu Cys Gln Asp Lys Asn Ile Leu Leu Tyr Asn Gln Arg Ser
 245 250 255
 Arg

<210> 24

<211> 257

<212> PRT

<213> Artificial Sequence

<220>

5 <223> Description of Artificial Sequence: Synthetic
Sequence

<400> 24

Met Thr Ala Ala Val Phe Phe Gly Cys Asp Phe Ile Ala Phe Gly Pro
1 5 10 15
10 Ala Leu Ala Glu Tyr Val Phe Thr Ile Ala Thr Glu Pro Phe Arg Ile
20 25 30
Ile Phe Leu Ile Ala Gly Ala Gly Phe Trp Leu Val Ser Leu Leu Ile
35 40 45
Ser His Leu Val Trp Phe Met Ala Arg Val Ile Lys Asp Asn Lys Asp
15 50 55 60
Gly Pro Thr Gln Lys Leu Leu Leu Ile Phe Gly Ala Phe Val Ser Met
65 70 75 80
Tyr Ile Gln Glu Met Phe Arg Phe Ala Asn Tyr Lys Leu Leu Lys Lys
85 90 95
20 Ala Ser Glu Gln Leu Lys Ser Ile Asn Pro Gly Glu Thr Arg Pro Ser
100 105 110
Met Arg Leu Leu Ala Tyr Val Thr Gly Leu Gly Phe Gly Ile Met Ser
115 120 125
Gly Trp Phe Ser Phe Val Asn Thr Leu Ser Asp Tyr Leu Gly Pro Gly
25 130 135 140
Thr Val Gly Ile His Ala Asp Ser Pro Gln Phe Phe Leu Tyr Ser Asp
145 150 155 160
Phe Met Thr Leu Val Ile Ile Leu Leu Glu Val Phe Trp Gly Ile Val
165 170 175
30 Phe Phe Asp Phe Cys Glu Lys Lys Lys Trp Gly Ile Leu Gly Ile Val
180 185 190
Leu Leu Thr His Leu Leu Val His Ala Gln Thr Phe Ile Ser Ser Tyr
195 200 205
Tyr Ile Ile Asn Leu Ala Ser Ala Phe Ile Ile Lys Val Leu Met Gly
35 210 215 220
Thr Trp Ala Phe Leu Leu Ala Gly Gly Ser Cys Arg Ser Leu Lys Met
225 230 235 240
Cys Leu Leu Cys Gln Asp Lys Asn Phe Asn Leu Tyr Asn Gln Arg Ser

245 250 255

Arg

<210> 25
<211> 925
5 <212> PRT
<213> Artificial Sequence
<220>
<223> Description of Artificial Sequence: Synthetic
Sequence

10 <400> 25
Gly Ala Thr Gly Ala Cys Thr Gly Cys Gly Gly Cys Cys Gly Thr Gly
1 5 10 15
Thr Thr Cys Thr Thr Cys Gly Gly Cys Thr Gly Cys Gly Cys Cys Thr
20 25 30
15 Thr Cys Ala Thr Thr Gly Cys Cys Thr Thr Cys Gly Gly Gly Cys Cys
35 40 45
Thr Gly Cys Gly Cys Thr Cys Gly Cys Cys Cys Thr Thr Thr Ala Thr
50 55 60
Gly Thr Cys Thr Thr Cys Ala Cys Cys Ala Thr Cys Gly Cys Cys Ala
20 65 70 75 80
Cys Cys Gly Ala Gly Cys Cys Gly Thr Thr Gly Cys Gly Thr Ala Thr
85 90 95
Cys Ala Thr Cys Thr Thr Cys Cys Thr Cys Ala Thr Cys Gly Cys Cys
100 105 110
25 Gly Gly Ala Gly Cys Thr Thr Thr Cys Thr Thr Cys Thr Gly Gly Thr
115 120 125
Thr Gly Gly Thr Gly Thr Cys Thr Cys Thr Ala Cys Thr Gly Ala Thr
130 135 140
Thr Thr Cys Gly Thr Cys Cys Cys Thr Thr Gly Thr Thr Thr Gly Gly
30 145 150 155 160
Thr Thr Cys Ala Thr Gly Gly Cys Ala Ala Gly Ala Gly Thr Cys Ala
165 170 175
Thr Thr Ala Thr Thr Gly Ala Cys Ala Ala Cys Ala Ala Ala Gly Ala
180 185 190
35 Thr Gly Gly Ala Cys Cys Ala Ala Cys Ala Cys Ala Gly Ala Ala Ala
195 200 205
Thr Ala Thr Cys Thr Gly Cys Thr Gly Ala Thr Cys Thr Thr Thr Gly
210 215 220

Gly Ala Gly Cys Gly Thr Thr Thr Gly Thr Cys Thr Cys Thr Gly Thr
 225 230 235 240
 Cys Thr Ala Thr Ala Thr Cys Cys Ala Ala Gly Ala Ala Ala Thr Gly
 245 250 255
 Thr Thr Cys Cys Gly Ala Thr Thr Thr Gly Cys Ala Thr Ala Thr Thr
 5 260 265 270
 Ala Thr Ala Ala Ala Cys Thr Cys Thr Thr Ala Ala Ala Ala Ala Ala
 275 280 285
 Ala Gly Cys Cys Ala Gly Thr Gly Ala Ala Gly Gly Thr Thr Thr Gly
 290 295 300
 10 Ala Ala Gly Ala Gly Thr Ala Thr Ala Ala Ala Cys Cys Cys Ala Gly
 305 310 315 320
 Gly Thr Gly Ala Gly Ala Cys Ala Gly Cys Ala Cys Cys Cys Thr Cys
 325 330 335
 Thr Ala Thr Gly Cys Gly Ala Cys Thr Gly Cys Thr Gly Gly Cys Cys
 15 340 345 350
 Thr Ala Thr Gly Thr Thr Thr Cys Thr Gly Gly Cys Thr Thr Gly Gly
 355 360 365
 Gly Cys Thr Thr Thr Gly Gly Ala Ala Thr Cys Ala Thr Gly Ala Gly
 370 375 380
 20 Thr Gly Gly Ala Gly Thr Ala Thr Thr Thr Thr Cys Cys Thr Thr Thr
 385 390 395 400
 Gly Thr Gly Ala Ala Thr Ala Cys Cys Cys Thr Ala Thr Cys Thr Gly
 405 410 415
 Ala Cys Thr Cys Cys Thr Thr Gly Gly Gly Gly Cys Cys Ala Gly Gly
 25 420 425 430
 Cys Ala Cys Ala Gly Thr Gly Gly Gly Cys Ala Thr Thr Cys Ala Thr
 435 440 445
 Gly Gly Ala Gly Ala Thr Thr Cys Thr Cys Cys Thr Cys Ala Ala Thr
 450 455 460
 30 Thr Cys Thr Thr Cys Cys Thr Thr Thr Ala Thr Thr Cys Ala Gly Cys
 465 470 475 480
 Thr Thr Thr Cys Ala Thr Gly Ala Cys Gly Cys Thr Gly Gly Thr Cys
 485 490 495
 Ala Thr Thr Ala Thr Cys Thr Thr Gly Cys Thr Gly Cys Ala Thr Gly
 35 500 505 510
 Thr Ala Thr Thr Cys Thr Gly Gly Gly Gly Cys Ala Thr Thr Gly Thr
 515 520 525
 Ala Thr Thr Thr Thr Thr Thr Gly Ala Thr Gly Gly Cys Thr Gly Thr

	530	535	540
	Gly Ala Gly Ala Ala Gly Ala Ala Ala Ala Gly Thr Gly Gly Gly		
	545	550	555 560
	Gly Cys Ala Thr Cys Cys Thr Cys Cys Thr Thr Ala Thr Cys Gly Thr		
	565	570	575
5	Thr Cys Thr Cys Cys Thr Gly Ala Cys Cys Cys Ala Cys Cys Thr Gly		
	580	585	590
	Cys Thr Gly Gly Thr Gly Thr Cys Ala Gly Cys Cys Cys Ala Gly Ala		
	595	600	605
	Cys Cys Thr Thr Cys Ala Thr Ala Ala Gly Thr Thr Cys Thr Thr Ala		
10	610	615	620
	Thr Thr Ala Thr Gly Gly Ala Ala Thr Ala Ala Cys Cys Thr Gly		
	625	630	635 640
	Gly Cys Gly Thr Cys Ala Gly Cys Ala Thr Thr Thr Ala Thr Ala Ala		
	645	650	655
15	Thr Cys Cys Thr Gly Gly Thr Gly Cys Thr Cys Ala Thr Gly Gly Gly		
	660	665	670
	Cys Ala Cys Cys Thr Gly Gly Gly Cys Ala Thr Thr Cys Thr Thr Ala		
	675	680	685
	Gly Cys Thr Gly Cys Gly Gly Gly Ala Gly Gly Cys Ala Gly Cys Thr		
20	690	695	700
	Gly Cys Cys Gly Ala Ala Gly Cys Cys Thr Gly Ala Ala Ala Cys Thr		
	705	710	715 720
	Cys Thr Gly Cys Cys Thr Gly Cys Thr Cys Thr Gly Cys Cys Ala Ala		
	725	730	735
25	Gly Ala Cys Ala Ala Gly Ala Ala Cys Thr Thr Thr Cys Thr Thr Cys		
	740	745	750
	Thr Thr Thr Ala Cys Ala Ala Cys Cys Ala Gly Cys Gly Cys Thr Cys		
	755	760	765
	Cys Ala Gly Ala Thr Ala Ala Cys Cys Thr Cys Ala Gly Gly Gly Ala		
30	770	775	780
	Ala Cys Cys Ala Gly Cys Ala Cys Thr Thr Cys Cys Cys Ala Ala Ala		
	785	790	795 800
	Cys Cys Gly Cys Ala Gly Ala Cys Thr Ala Cys Ala Thr Cys Thr Thr		
	805	810	815
35	Thr Ala Gly Ala Gly Gly Ala Ala Gly Cys Ala Cys Ala Ala Cys Thr		
	820	825	830
	Gly Thr Gly Cys Cys Thr Thr Thr Thr Cys Thr Gly Ala Ala Ala		
	835	840	845

Ala Thr Cys Cys Cys Thr Thr Thr Thr Thr Cys Thr Gly Gly Thr Gly
 850 855 860
 Gly Ala Ala Thr Thr Gly Ala Gly Ala Ala Ala Gly Ala Ala Ala Thr
 865 870 875 880
 Ala Ala Ala Ala Cys Thr Ala Thr Gly Cys Ala Gly Ala Thr Ala Thr
 5 885 890 895
 Gly Cys Gly Thr Thr Cys Cys Ala Ala Ala Ala Ala Ala Ala Ala
 900 905 910
 Ala Ala Ala Ala Ala Ala Ala Ala Ala Ala Ala Ala Ala
 915 920 925

10
 <210> 26
 <211> 925
 <212> DNA
 <213> human

15
 <400> 26
 gatgactgcg gccgtgttct tcggtgcgc cttcattgcc ttcgggcctg cgctcgccct 60
 ttatgtcttc accatcgcca ccgagccgtt gcgtatcatc ttcctcatcg ccggagcttt 120
 cttctggttg gtgtctctac tgatttcgtc ccttgtttgg ttcattggca gagtcattat 180
 tgacaacaaa gatggaccaa cacagaaata tctgctgac tttggagcgt ttgtctctgt 240
 20 ctatatccaa gaaatgttcc gatttgcata ttataaactc ttaaaaaaag ccagtgaagg 300
 tttgaagagt ataaaccag gtgagacagc accctctatg cgactgctgg cctatgtttc 360
 tggtctgggc tttggaatca tgagtggagt attttccttt gtgaataccc tatctgactc 420
 cttggggcca ggcacagtgg gcattcatgg agattctcct caattcttcc tttattcagc 480
 tttcatgacg ctgggtcatta tcttgcctga tgtattctgg ggcattgtat tttttgatgg 540
 25 ctgtgagaag aaaaagtggg gcacctcctc tatcgttctc ctgacccacc tgctggtgtc 600
 agcccagacc ttcataagtt cttattatgg aataaacctg gcgtcagcat ttataatcct 660
 ggtgctcatg ggcacctggg cattcttagc tgcgggaggc agctgccgaa gcctgaaact 720
 ctgcctgctc tgccaagaca agaactttct tctttacaac cagcgtcca gataacctca 780
 gggaaccagc acttcccaa cgcagacta catctttaga ggaagcaca ctgtgccttt 840
 30 ttctgaaaat ccctttttct ggtggaattg agaaagaaat aaaactatgc agatatgctg 900
 tccaaaaaaa aaaaaaaaaa aaaaa 925

<210> 27
 <211> 925
 35 <212> DNA
 <213> Artificial Sequence
 <220>
 <223> Description of Artificial Sequence: Synthetic

Sequence

<400> 27

gatgacagcg gccgttttct tcggetgccc cttgattgcc ttagggcctg ctctcgccct 60
 ctatgtcttg accatcgcaa ccgagccctt gcgtatcatc ttcctgatcg ccggagcttt 120
 cttttgggtg gtctctctac tgatttcgtc acttgtttgt ttcattggcca gagtcattgat 180
 5 tgacaaaaaa gatgggtccaa cacacaaata tctgctgac ttaggagcgt ttgtctctgt 240
 ctatatccag gaaatgttac gatttgctta ttataacctc ttaaagaaag ccagagaagg 300
 ttttaagagt atcaaccag gggagacagc accctctatt cgactgctcg cctatgtgtc 360
 tggetttaggc tttggatca tgagcggagt attgtccttt gtaaataccc tgtctgactc 420
 cttggggccg ggcacagtag gcattcaggg agattccctt caattgttcc tttaatcagc 480
 10 ttttatgacg ctctcatta tgttgcgtca agtattctgt ggcattgtct tttttgaggg 540
 ctgtgaaaag aaaaattggg gcacccctct tatgggtctc ctaaccacc ttctgggtgtc 600
 cgcccagacg ttcataagat cttattatgg aataaacctg gcgtcggcat ttataatcct 660
 ggttctcatg ggcacctggg cgttcttagc agcgggaggt agctgccga gcctgaagct 720
 ctgcctactc tgccatgaca agaactttct tctgtacaac caacgctcca gttaacctca 780
 15 cggaaccagg acttccaaa ccgcagatta catcttcaga ggaaggacaa ctgtaccttt 840
 ttctgaaaat ccttttttct ggtggaattg agaaagaaat aaaactatcc agatatgggt 900
 tccaaaaaaa aaaaataaaa aaac 925

<210> 28

20 <211> 925

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic

25 Sequence

<400> 28

gatgactgcg gccgtcttct tcgggtgccc ctttaattgcc tttgggcctg cctcgcct 60
 gtatgtctta accatcgcta ccgagccctt gcgtatgac ttcctaactg ccggtgcttt 120
 cttctgggtg gtgtctctac taatttcgtc tcttggttgc ttcattggca gagtcataat 180
 30 tgacaataaa gatggcccaa cacagaaata tctactgac tttggagcgt tcgtctctgt 240
 gtatatccaa gaaatgttct gatttgccta ttataagctc ttaaaaaaag ccagtgaagg 300
 tttcaagagt atgaaccag gagagacagc tccctctatc cgactgctgg cctatgtatc 360
 tggctttggc tttggcatca tgaggggagt attatccttt gtttaataccc tctctgactc 420
 gttggggcca ggcacagttg gcattcacgg agattcgctt caattattcc tttattcagc 480
 35 tttcatgacg ctggctatta tattgctgca tgtattctgc ggcattgtgt tttttgaagg 540
 ctgtgataag aaaaactggg gcattgctct tatagttctc cttaccacc tctgggtgtc 600
 ggcccagaca ttcataagtt cttattacgg aataaagctg gcgtcagcat ttattatcct 660
 ggtcctcatg gggacctggg cattcttagc tgcgggaggg agctgccgga gcctgaaact 720

ctgccttctc tgccacgaca agaagtttct tctatacaac catcgctcca gctaacctca 780
 gggaaccaga acttcccata ccgcagacta catcttgaga ggaagaacaa ctgttccttt 840
 ttccgaaaat ccgtttttct gatggaattg tgaaagaaac aaaactatgc agatatgagt 900
 tccaaataaa aaaaacaaaa aaaag 925

5 <210> 29

<211> 925

<212> DNA

<213> Artificial Sequence

<220>

10 <223> Description of Artificial Sequence: Synthetic
 Sequence

<400> 29

gatgaccgcg gccgtgttct tcggatgcgc ctttattgcc ttggggcctg cgctcgccct 60
 atatgtcttt accatcgcca ccgagccgtt gcgtataatc ttccctatcg ccggcgcttt 120
 15 cttgtggttg gtatctctac ttatttcgtc ccttggttgg ttcattggca gagtcattat 180
 tgacaacaaa gatgggcca caaaaaata tcttctgac ttgggagcgt tggctctctgt 240
 atatatccat gaaatgttcc gatttgcgta ttataaactc ttaaataaag ccagcgaagg 300
 tttgaagagt ataaaccag gtgagacagc ccctctatg cgactgctag cctatgtttc 360
 tggcttcggc tttgggatca tgagaggagt attttccttt gtcaataccc tgtctgactc 420
 20 attggggcct ggcacagtcg gcattcaggg agattcacct caatttttcc tttactcagc 480
 tttgatgacg ctatgcatta ttttgcgca cgtattctgg ggcattgtat tttttgatgg 540
 ctgtgacaag aaaaagtggg gcatactcct tattgttctc ctcacccacc tgctggtgtc 600
 agcccagact ttcataagct cttattaggg aataaaaactg gcgtctgcat ttatcatcct 660
 ggtgctcatg ggaacctggg ctttcttagc cgcgggaggg agctgccgaa gcctgaatct 720
 25 ctgcctcttc tgccaggaca agaaatttct tctttacaac caccgctcca ggtaacctca 780
 aggaaccagt acttcccaca ccgcagagta catcttaaga ggaagtacaa ctgtcccttt 840
 ttccgaaaat ccatttttct gttggaattg cgaaagaaag aaaactatac agatatgtgt 900
 tccaaacaaa aaaaagaaaa aaaaa 925

30 <210> 30

<211> 925

<212> DNA

<213> Artificial Sequence

<220>

35 <223> Description of Artificial Sequence: Synthetic
 Sequence

<400> 30

gatgacggcg gccgtattct tcggttgcgc cttcattgcc ttggggcctg cactcgccct 60

ttatgtcttc accatcgca cagagccatt gcgtattatc ttcctcatcg ccggggcttt 120
 cttatggttg gtttctctac tcatttcgtc gcttggttga ttcattggcta gagtcatcat 180
 tgacaagaaa gatggaccaa cacataaata tctcctgac ttgggagcgt tagtctctgt 240
 ttatatccac gaaatgttgc gatttgcata ttataatctc ttaaacaag ccagggaagg 300
 tttaaagagt attaaccag gcgagacagc gccctctata cgactgcttg cctatgtctc 360
 5 tggcttgggc tttggaatca tgagtggagt attctccttt gtgaataccc tatctgactc 420
 tttggggccc ggcacagtgg gcattcaagg agattctcct caattcttcc tttagtcagc 480
 tttaatgacg cttgtcatta tcttgctgca ggtattctga ggcattgttt tttttgacgg 540
 ctgtgagaag aaaaaatggg gcattctcct tatcgttctc ctgaccacc tactggtgtc 600
 tgcccagacc ttcataaggt cttattaagg aataaatctg gcgtccgcat ttatgacct 660
 10 ggtactcatg ggtacctggg ccttcttagc ggcgggagga agctgccgta gcctgaacct 720
 ctgcctgtc tgccaagaca agaattttct tctctacaac cagcgtcca gataacctca 780
 tggaaccagc acttcccaga ccgcagaata catctttaga ggaagcaca ctgtgccttt 840
 ttcagaaaat ccttttttct gctggaattg ggaaagaaaa aaaactattc agatatgcgt 900
 tccaaagaaa aaaaaaaaaa aaat 925

15

<210> 31

<211> 925

<212> DNA

<213> Artificial Sequence

20

<220>

<223> Description of Artificial Sequence: Synthetic
 Sequence

<400> 31

gatgactgca gccgtgtttt tcggctgcgc cttcatggcc ttcggacctg cgcttgccct 60
 25 ttacgtcttc acgatcgcca cagagccgtt tcgtatcatc ttcctcatgg ccggagcatt 120
 cttctgtttg gtgtccctac tgatgtcgtc cctagtttgg tttatggcaa gcgtcattat 180
 ggacaacaaa gatggacctc cacagaacta tctgctgac tttggagcgt ttgtttctgt 240
 ctacatccaa gagatgttcc gatttgcata ttataaactc ttaaaaaagg ccagtgaagg 300
 tttgaatagt ataaaccag gtgagacagc accatctatg cgtctgctgg cctatgtttc 360
 30 gggcttggga tttggaatta tgagtggcgt attttcgttt gtgaaaacc tatctgactc 420
 cttegggcca gggacagtgg gaattcatgg tgattctccc caattcttgc tttattcagc 480
 ttccattacg ctggtcatta tcttgctgca tgtattctgg ggtattgtat tctttgatgg 540
 gtgtgagaaa aaaaagtgtg gcacctcct tatcgtgctc ctgacacacc tgcttggtgc 600
 agcccagacc ttgataagtt catattatgg tataaacctc gcgtcagcgt ttataatact 660
 35 ggtgcttatg ggcacctggg cattgttagc tgcaggaggc agttgccgaa gcctgaaact 720
 gtgcctgcta tgccaagata agaacttctc tctttagaac cagcgatcca gatatcctca 780
 gggcaccagc acgtcccaaa cagcagacta tatctttagc ggaagcacga ctgtgccatt 840
 ttctgataat cccttcttct ggtggaattg agaaagaaat aatactatgc acatagcgt 900

gccaaaaaaa aaaaaaata aaaaa

925

<210> 32

<211> 925

<212> DNA

5 <213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
Sequence

<400> 32

10 gatgactgct gccgtgttct tcggctgggc cttcatagcc ttcggtcctg cgctcgccct 60
 ttaggtcttc acaatcgcca ctgagccgtt ccgtatcatg ttctcatag cggagcttt 120
 cttctgcttg gtgtcgctac tgatatcgtc ccttgtttg ttcatggcaa gggcattat 180
 agacaacaat gatggaccca cacagaagta tctgctaata tttggtgcgt ttgtctctgt 240
 ctagatccaa gaaatgttcc gttttgcata ctataaactg ttaaaaaaag ccagtgatgg 300
 15 tttgaacagt ataaagccag gtgaaacagc accttctatg cgctgctgg cgtatgttct 360
 aggcttgggt tttggaatca tgagtggggg attttcattt gtgaataccc tatccgactc 420
 cttggggcca ggaacagtgg gtattcatgg cgattctccg caattcttac tttattctgc 480
 tttcatcacg ctgggtgatta tcttactgca tgttttctgg ggcattgtat tgtttgatgg 540
 atgtgagaat aaaaagtgcg gcatcctgct tatcgtactc ctgactcacc tgctcgtgtc 600
 20 agcgcagacc ttaataagtt cttattatgg cataaacctg gcgtcagcat ttataattct 660
 ggtgctcatg ggcaagtggg cattattagc tgctggaggc agctgccgaa ggctgaaact 720
 atgcctgctt tgccaagaca agaacttgct tctttaaac cagcgttcca gataccctca 780
 ggggaccagc acatcccaaa ctgcagacta catctttagg ggaagcacia ctgtgccttt 840
 ttctgacaat cccttgttct ggtgaaattg agatagaaat aacactatgc agatatgcgt 900
 25 accaaaaaat aaaaaaaca aaaaa 925

<210> 33

<211> 925

<212> DNA

30 <213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
Sequence

<400> 33

35 gatgactgcc gccgtgttgt tcggctgagc cttcattgcc ttccggccctg cgctggccct 60
 ttaagtcttc actatcgcca ccgagccgtt gcgtatcata ttctcattg ccggagcctt 120
 cttctggttg gtgtcactac tgatttcgtc cctcgtttg ttgatggcaa gagtcattat 180
 tgacaacaac gatggaccga cacagaaata tctgcttata tttggcgct ttgtgtctgt 240

ctaaattccaa gatattgttcc gctttgcata gtataaacta ttaaaaaatg ccagtgcagg 300
 tttgaagagt ataaaaccag gtgatacagc accctctatg cggctgctgg catatgtttc 360
 tggcttgggc tttggaatga tgagtggagt attttctttt gtgaacaccc tatcggactc 420
 cttagggcca ggtacagtgg gcattcatgg ggattctcca caattctttc tttattccgc 480
 tttcatgacg ctggtaatta tctttctgca tgtcttctgg gggattgtat tatttgatgg 540
 5 tttgtgagaac aaaaagtggg gcatactact tatcgttctc ctgacccacc tgcgtggtgc 600
 agcacagacc tttataagtt cctattatgg gataaaccta gcgtcagctt ttataatcct 660
 ggtgctgatg ggcacatggg catttttagc tgccggaggc aggtgccgaa gactgaaact 720
 ttgctgctc tgccaagaga agaacttact tctttataac cagcgtccca gatagcctca 780
 gggaaccagc acttcccaaa ccgcagacta gatctttaga ggaagcacta ctgtgccctt 840
 10 ttctgagaat cccttattct ggtgtaattg agacagaaat aagactatgc aaatatgcgt 900
 tccaaaaaac aaaaaaaga aaaaa 925

<210> 34

<211> 925

15 <212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
Sequence

20 <400> 34

gatgactgcg gccgtgttat tcggctgtgc cttcatcgcc ttccggcctg cgctagccct 60
 ttatgtcttc accatcgcca cggagccgtt acgtatcatt ttctctatcg ccggagcgtt 120
 cttctgattg gtgtctctac tgatctcgtc cctggtttgg ttaatggcaa gtgtcattat 180
 cgacaacaag gatggaccaa cacagaatta tetgctcacc tttggggcgt ttgtatctgt 240
 25 ctatatccaa gacatgttcc ggtttgcata atataaactt ttaaaaaacg ccagtgcagg 300
 tttgaaaagt ataaatccag gtgacacagc accgtctatg cgactgctgg cttatgtttc 360
 cggcttgggg tttggaataa tgagtgggtg attttctttt gtgaagaccc tatcagactc 420
 ctttgggcca ggcacagtgg ggattcatgg agattctcct caattcttcc tttattccgc 480
 tttcataacg ctggttatta tcttctgca tgtgttctgg ggaattgtat tttttgatgg 540
 30 ctgtgagaag aaaaagtgag gcatacttct tatcgtctc ctgacgcacc tgctagtgtc 600
 agctcagacc ttcataagtt cgtattatgg aataaacctt gcgtcagcct ttataatgct 660
 ggtgctaata ggcaacttgg cattcttagc tgccggaggc agatgccgaa gtctgaaact 720
 ctgctgctg tgccaagaaa agaactttct tctttacaac cagcgtccca gataacctca 780
 ggtaccagc acctcccaaa ccgcagacta aatctttagt ggaagcacca ctgtgccgtt 840
 35 ttctgaaaat ccctttttct ggtgcaattg agagagaaat aaaactatgc atatatgcgt 900
 cccaaaaaag aaaaaaaaaa aaaaa 925

<210> 35

<211> 925

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic

5

Sequence

<400> 35

gatgactgcg gcagtgttct ttggctgccc cttcattgcg ttcgggccag cgctcgctct 60
 ttatgtcttc accatggcca ccgaaccgtt gcgtatcatc ttcctcatcg cgggagcttt 120
 attctggttt gtgtctctcc tgatttcgtc ccttgatagg ttcattgcaa gagtcattat 180
 10 tgagaacaaa gaaggaccaa ctcaaaaata cctgctgatg tttggagcat ttgtctctgt 240
 ctatatccaa gaaatgttcc gattagcata ttataaactc ttcaaaaaag cgagtgaagg 300
 attgaagagt ataaaccccg gtgagacggc accctcaatg cgacttctgg cctacgtttc 360
 tgggttgggc ttaggaatca ttagtggagt cttttccttg gtgaatacac tatctgattc 420
 cttgggcccc ggacacgttg gcatacatgg agattctcct cacttcttcc tgtattcagc 480
 15 attcatgact ctggatcatc tcttgctgca tgtattatgg ggcatgtgat ttttcgatgg 540
 ctgggagaag aaaaagtggg gtatcctcct catcgttctg ctgacccaac tgctggtttc 600
 agcccacacc ttcattgagt cttaatatgg aattaacctg gcctcagcat tgataatcct 660
 agtgctcatt ggacacgtcg cattcttggc tgcgggaggg agctgtcgaa gcctcaaaact 720
 ctggctgctc tgacaagaca ataactttct cctttacaag cagcgctcaa gataacctca 780
 20 gggaaccagc acttcgcaaa ccgcagacta cattttttaga ggcagcacia cgggtgccttt 840
 atctgaaaat ccctttttct ggtggaagtg agaaagaaat aaaactatgc agatctgcgt 900
 tccgaaaaaa aaaaaaaaaa ataaa 925

<210> 36

25

<211> 925

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic

30

Sequence

<400> 36

gatgactgcg gctgtgttct tcggctgccc gttcattgca ttcgggcctg cgctcgccct 60
 ttatgtgttc accatagcca ccgatccgtt gcgcatcatc ttgctcatcg caggagcttt 120
 tttctggttc gtgtctctgc tgatttcac ccttgtttgg ttcacgcaa gagtgattat 180
 35 tgaaaacaaa gatggaccaa ccagaaaata gctgctgata tttggagctt ttgtctccgt 240
 ctatatgcaa gaaatattcc gatttgcata ttacaaactc ttgaaaaaag caagtgaagg 300
 tttgaagagc ataaacccgg gtgagacagc accctctatg cgactcctgg cctagggttc 360
 tggattgggc tttggaatca tcagtggagt gttttcctta gtgaatactc tatctgactc 420

cttggggcca ggcacagtgg gcattcatgg agactctcct cagttcttcc tatattcagc 480
 tttcatgacc ctggatcatga tcttgctaca tgtatttttg ggcatcgat ttttggatgg 540
 ctgagagaag aataagtggg gcacccctct gatcggttcta ctgaccatc tgctgggtctc 600
 agcccagacc ttcataagtt cttattatgg aatcaacctg gcgtcagcat taataatcct 660
 tgtgtctatc ggcacctggg cattcttagc tgcgggtggc agctgccgaa gcctgaaact 720
 5 ctgactgctc tgtcaagaca acaactttct gctttacaaa cagcgctcta gataacccca 780
 gggaacgagc acttcacaaa ccgctgacta catctttaga gggagcacia cagtgccttt 840
 ttctgaaaac ccctttttgt ggtggaaatg agaaagtaat aaaacatgc agatgtgcgt 900
 tccaaaaaaa aataaaaaaa acaaaa 925

10 <210> 37

<211> 925

<212> DNA

<213> Artificial Sequence

<220>

15 <223> Description of Artificial Sequence: Synthetic
Sequence

<400> 37

gatgactgcg gccgtgttct tgggctgccc attcattgct ttccggcccg cgctcgcgct 60
 ttatgtattc accattgccg ccgaccggtt gggatcatc ttactcatcg ctggagcttt 120
 20 cttctgggtg gtgtctctac tgatttcttc ccttgtctgg ttcattggca gagtaattat 180
 tgataacaaa gacggaccaa cgcagaaata actgctgatt tttggagcct ttgtctcggt 240
 ctatatacaa gaaattttcc gattcgcata ttagaaactc ttaaaaaaag ctagtgaagg 300
 cttgaagagg ataaaccag gtgagactgc accctccatg cgactgctgg cctaagtttc 360
 tggtttgggc ttccgaatca tgagtggagt attttcttt gtgaataccc tatctgagtc 420
 25 cttgggacca ggcactgtgg gcacccatgg agagtctcct caattcttcc tttattcagc 480
 cttcatgacg ctggatcataa tcttgcttca tgtattctgg ggcatggat ttttagatgg 540
 ctgtgagaag aacaagtggg ggatcctcct aatcggtctt ctgaccacc tgctgggtgc 600
 agcccaaacc ttcattagtt cttactatgg aatgaacctg gcacagcat ttataatcct 660
 cgtgtcatg ggcacctgag cattctttgc tgcggggcggc agctggcgaa gcctaaaact 720
 30 ctgtctgctc tgccaagaca agaactttct actttacaat cagcgctcca gataaccgca 780
 gggaacaagc acttctcaaa ccgccgacta catgtttaga ggaagcacia ctgtgccttt 840
 ctctgaaaag ccctttttat ggtggaattg agaaagcaat aaaacgatgc agatatgcgt 900
 tcctaaaaaa aacaaaaaaa agaaa 925

35 <210> 38

<211> 925

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
Sequence

<400> 38

```
gatgactgcg gcggtgttct taggctgcgc ttccattgcc ttggggccgg cgctcgact 60
5  ttatgttttc accatcgcca ccgagccgtt gcgaatcacc ttctcatcg ccggagcttt 120
   gttctggtta gtgtctcttc tgatttcctc ccttggtggg ttcatagcaa gagttattat 180
   tgacaacaaa gagggaccaa cacagaaata tctgctgac tttggagcgt ttgtctcagt 240
   ctatattcaa gaaatcttcc gattggcata ttaaaaactc tttaaaaaag ccagtgaagg 300
   gttgaagaga ataaaccctg gtgagaccgc accctcgatg cgactactgg cctatgtttc 360
10  tggcttgggc ttgggaatca taagtggagt ttttctcttc gtgaatacgc tatctgaatc 420
   cttgggtcca ggcaccgtgg gcatgcatgg agaattctct catttcttcc tctattcagc 480
   gttcatgaca ctggctatta tcttgctcca tgtattgtgg ggcatagtat tttttgatgg 540
   ctgcgagaag aagaagtggg gaatcctcct tatcgttctc ctgaccagc tgctgggtatc 600
   agcccatacc ttcattcagtt cttagtatgg aataaacctg gcttcagcat tcataatcct 660
15  ggtgctcata ggcacctgtg cattcttcgc tgcggggggc agctgacgaa gccttaaact 720
   ctgcctgctc tggcaagaca aaaactttct tctttacaac cagcgctcga gataaccaca 780
   gggaactagc acttcccaaa ccgcggacta catatttaga ggtagcacia ccgtgccttt 840
   gtctgaaaaa cccttttttt ggtggaactg agaaaggaat aaaacaatgc agatttgcgt 900
   tcccaaaaaa aagaaaaaaa aaaaa 925
```